# Wetland Assessment at Homestead Bay, Queenstown

Contract Report No. 7483a

Providing outstanding ecological services to sustain and improve our environments





## Wetland Assessment at Homestead Bay, Queenstown

#### Contract Report No. 7483a

February 2025

#### **Project Team:**

Erin Gallagher - Report author and field survey Kelvin Lloyd - Report review

#### Prepared for:

RCL Homestead Bay Limited Level 1, 13 Camp Street Queenstown 9300

Reviewed and approved for release by:



Kelvin Lloyd Senior Principal Ecologist Wildland Consultants Ltd 14/02/2025

Cite this report as follows:

Wildland Consultants (2025). *Wetland Assessment at Homestead Bay, Queenstown.* Wildland Consultants Contract Report No. 7483a. Prepared for RCL Homestead Bay Limited. 56 pp.

#### **Dunedin Office**

764 Cumberland Street, Dunedin, Ph 03-477-2096

#### Head Office

99 Sala Street, PO Box 7137, Te Ngae, Rotorua Ph 07-343-9017 Fax 07-343-9018 Email: rotorua@wildlands.co.nz

www.wildlands.co.nz



## **Contents**

1.0	Introduction	3
2.0	Project Scope	3
3.0	Relevant Legalisation	3
3.1	Wetland definitions	3
4.0	Methods	5
4.1	Desktop analysis and considerations	5
4.2	Field survey	5
4.3	Wetland assessment	6
5.0	Potential wetland areas	7
5.1	Overview	7
5.2	Wet Area 1	9
5.3	Wet Area 2	10
5.4	Wet Area 3	11
5.5	Wet Area 4	11
5.6	Wet Area 5	12
5.7	Wet Area 6	12
5.8	Wet Area 7	13
5.9	Wet Area 8	13
5.10	Wet Area 9	13
6.0	Natural inland wetlands	14
6.1	Overview	14
7.0	Summary	20
Ackno	owledgments	20
Refer	rences	20
Appe	endix 1	22
Plant	species recorded and their wetland indicator status	22
Appe	endix 2	24
	thly rainfall data for 2024, 2025 and historical averages	24
Appe	endix 3	25
	photographs	25
Appe	endix 4	29
Wetla	and delineation plot sheets	29

#### © Wildland Consultants Ltd 2025

This report has been produced by Wildland Consultants Ltd for RCL Homestead Bay Limited. All copyright in this report is the property of Wildland Consultants Ltd and any unauthorised publication, reproduction, or adaptation of this report, including entry into generative artificial intelligence, is a breach of that copyright.





### 1.0 Introduction

Remarkable Planning, on behalf of RCL Henry Downs Limited, is preparing a Fast Track resource consent application for a residential subdivision on approximately 41 hectares of rural land at Homestead Bay, just south of Jacks Point in Queenstown, Otago (Figure 1). The gently sloping site has been farmed for many decades and is largely vegetated in exotic pasture, but also contains small areas of tūmatakuru/matagouri (*Discaria toumatou*) shrublands, gullies with ephemeral streams, and possibly small natural inland wetlands. Previous aquatic and terrestrial ecology assessments have been undertaken for the site, but further assessments of natural inland wetlands are now also required.

RCL Homestead Bay Limited has commissioned Wildland Consultants Ltd (Wildlands) to identify any wetlands present and determine whether these would be exempt from the definition of a natural inland wetland under Clause 3.21(e) of the National Policy Statement for Freshwater Management (Ministry for the Environment 2020b).

## 2.0 Project Scope

The scope of this project includes:

- Identification of any wetlands within the proposed development.
- Determine if any wetlands present meet the definition of a natural inland wetland under Clause 3.21 of the National Policy Statement for Freshwater Management (NPS-FM; October 2024 amendments).

## 3.0 Relevant Legalisation

#### 3.1 Wetland definitions

Wetlands have been defined in the Resource Management Act (RMA, 'the Act'), as outlined below.

Wetland – permanently or intermittently wet areas, shallow water, and land/water margins that support a natural ecosystem of plants and animals that are adapted to wet conditions, including within the coastal marine area.

The National Policy Statement for Freshwater Management (NPS-FW) defines 'natural inland wetlands' as outlined below.

Natural inland wetland means a wetland (as defined in the Act) that is not:

- (a) in the coastal marine area; or
- (b) a deliberately constructed wetland, other than a wetland constructed to offset impacts on, or to restore, an existing or former natural inland wetland; or
- (c) a wetland that has developed in or around a deliberately constructed water body, since the construction of the water body; or
- (d) a geothermal wetland; or







#### (e) a wetland that:

(i) is within an area of pasture used for grazing; and

(ii) has vegetation cover comprising more than 50% exotic pasture species (as identified in the National List of Exotic Pasture Species using the Pasture Exclusion Assessment Methodology (see clause 1.8)); unless

(iii) the wetland is a location of a habitat of a threatened species identified under clause 3.8 of this National Policy Statement, in which case the exclusion in (e) does not apply.

According to this definition, the pre-requisite for an area to be classified as a natural inland wetland is for the area to meet the wetland definition under the RMA, which requires both suitable hydrological conditions and presence of plants that are adapted to wet conditions, but which must not meet any of the exceptions listed above.

## 4.0 Methods

#### 4.1 Desktop analysis and considerations

An initial search was undertaken on Google Earth to identify potential wetland areas to survey.

The hydrological guidance for accurate wetland delineation recommends that site inspections should be undertaken under 'normal' hydrological conditions and within the growing season for plants in the relevant region. Normal hydrological conditions require 'normal' expected rainfall for the two to three months prior to the site inspection, and for the site inspection to not be undertaken following a period of heavy rain.

The growing season for the lower areas around Queenstown starts in September and ends in May (Ministry for the Environment 2021). This survey was undertaken within the appropriate local growing season (January and February 2025).

Rainfall for January 2025 (26.8 millimetres) was significantly lower than historical average (58.1 millimetres). Rainfall for December 2024 was higher (69.8 millimetres) than the historical average of 56.1 millimetres. Rainfall for November 2024 (57.8 millimetres) was very slightly higher than the historical average of 56.5 millimetres) (Appendix 2, Metservice 2025). The dryer than normal conditions of January 2025 were taken into consideration when assessing all potential wetlands.

The soil temperature was not measured during this assessment. Soil temperature is taken to enable the determination of the growing season. However, the growing season was determined by using the guidelines in the Wetland Hydrology Tool (Ministry for the Environment 2021).

#### 4.2 Field survey

Site investigations were undertaken on 30 January, 3 and 4 February 2025 to assess potential wetlands. Wetland vegetation types were classified and described following the structural classes outlined in Atkinson (1985). Wetland types were classified and described following the classification outlined in Johnson and Gerbeaux (2004).

Another site visit was undertaken on 3 September 2025. The purpose of the site visit was to investigate another potential wetland within an ephemeral stream and to perform a hydric soil and hydrology



assessment for a potential wetland that was within the 'Wet Area 1' of the wetland assessment undertaken in February 2025 (Wildland Consultants 2025).

#### 4.3 Wetland assessment

#### Part 1: Assess wetland status under the RMA

To define whether a wetland meets the RMA definition of a wetland, the Part 1 assessment within the defining 'natural wetlands' and 'natural inland wetlands' guidance document needs to be undertaken (Ministry for the Environment 2021). This assessment can also help define the extent of any wetland present.

The New Zealand vegetation tool for wetland delineation (Clarkson 2013) has become the standard methodology to assess the presence of plants adapted to wet conditions. This methodology classifies all plant species recorded in wetlands into five categories.

- OBL: Obligate. Almost always is a hydrophyte, rarely in uplands (estimated probability >99% occurrence in wetlands).
- FACW: Facultative Wetland. Usually is a hydrophyte but occasionally found in uplands (estimated probability 67-99% occurrence in wetlands).
- FAC: Facultative. Commonly occurs as either a hydrophyte or non-hydrophyte (estimate probability 34-66% occurrence in wetlands).
- FACU: Facultative Upland. Occasionally is a hydrophyte but usually occurs in uplands (estimated probability 1-33% occurrence in wetlands).
- UPL: Obligate Upland. Rarely is a hydrophyte, almost always in uplands (estimated probability <1% occurrence in wetlands).

Species that are classed as OBL, FACW, or FAC are considered hydrophytic and generally indicative of wetland habitat. The relative dominance of each species and corresponding classification can therefore determine whether an area should be defined as a wetland. In the instance, that a plant species present does not have a wetland indicator status, it given the UPL status (Clarkson *et al.* 2021).

In accordance with the methods described in the wetland delineation protocols (Clarkson 2021, Ministry for the Environment 2020, Ministry for the Environment 2021, Ministry for the Environment 2022 and Fraser *et al.* 2013), in areas of potential wetland, the following methods were applied:

- Firstly, the Rapid wetland test was completed. For this test to confirm the area as a wetland, all dominant species must be either OBL or FACW species. If the Rapid Wetland test failed, additional hydrophytic vegetation tests are required.
- Two tests are required for the hydrophytic vegetation determination (Dominance test and Prevalence index). Representative plots (2 metre × 2 metre for herb strata, 5 metre radius circular plot for shrub strata and 10 metre circular plot for tree/forest strata) where established in different vegetation types and geomorphic positions across the site. In each plot, the species in each stratum were identified and percent cover estimated (i.e. tree, sapling/shrub, herb). Species hydrophytic categories were taken from Manaaki Whenua (2021) and the dominant species were noted. For the Dominance test to confirm the area as a wetland, >50% of the dominant species must be OBL, FACW or FAC and all/most dominant species must not be FAC.
  - For the Prevalence Index (PI) test, a plot-based algorithm derived from the unique combination of OBL—UPL plants and their cover is calculated. The vegetation is hydrophytic (wetland) if PI ≤



- 3.0, but values around 3.0 can also be considered wetlands when other wetland indicators indicate wetland presence.
- If the Dominance, and Prevalence tests failed to identify the area as a wetland, then indicators of hydric soils and wetland hydrology were taken to determine if there was wetland hydrology present. If one of the hydrophytic vegetation tests passed and the other failed or if the result was uncertain (PI = 3.0 or Dominance test = 50%), further assessment is required.
- Methods for the soil assessment to determine hydric soils were taken from Fraser et al. (2013).
   The simple flow key (Figure 19) was followed to determine hydric soil features or other soil (or uncertain soil).
- Methods for the hydrology assessment were taken from Ministry for the Environment (2021). Wetland hydrology is determined by the presence or absence of hydrological indicators. Wetland hydrology indicators are assembled into four groups: 1) observation of flooding or groundwater; 2) evidence of flooding or ponding; 3) soil saturation; 4) landscape, vegetation and soil observations. Group 1 are primary indicators and Groups 2 to 4 have a mix of primary and secondary indicators. The presence of one primary indicator, or two secondary indicators, confirms the presence of a wetland.
- If the hydric soils and wetland hydrology tests are passed, then the definition of a wetland is met for the site under the Resource Management Act (RMA), but may not pass under the National Policy Statement for Freshwater Management (NPS-FM). If the hydric soils fail and the wetland hydrology pass, then this is also defined as a wetland. If the hydric soils pass and the wetland hydrology fail, then the area is a drained wetland or atypical environmental conditions are present. A site assessment is needed to determine the status in the latter case.

## <u>Part 2: Assessing whether a wetland is a 'natural wetland' or 'natural inland wetland' under the NPS-FM</u>

Once a wetland has been defined under the RMA, further assessment is needed to define whether a 'wetland' is a 'natural wetland' or a 'natural inland wetland'. The Part 2 assessment was followed within the defining 'natural wetlands' and 'natural inland wetlands' guidance document (Ministry for the Environment 2021). Exotic pasture species are noted from the National list of exotic pasture species document by Cosgrove *et al.* 2022.

### 5.0 Potential wetland areas

#### 5.1 Overview

RCL Homestead Bay Limited and Remarkable Planning identified two areas of potential wetlands (Wet Area 1 and 2). An initial and brief google earth analysis was undertaken at the proposal phase of this project and another possible five wetlands were identified (Wet Areas 3-5). Three of these potential wetlands are likely to be ephemeral wetlands due to their shape and geographic position which can be seen aerially on google earth. During the field survey, another area was also suspected to be a wetland (Wet Area 6). A total of nine areas were assessed for potential wetlands (Figure 1 and Table 1). Plant species recorded within the Wet Areas are listed in Appendix 1.

All potential wetland areas identified are currently within grazed pasture (improved pasture). The clause in the NPS-FM that concludes that a wetland that 'is within an area of pasture used for grazing' is not a natural inland wetland does not apply to these potential wetland areas as it is being proposed for residential development.



Table 1 - Wetland delineation results summary for potential wet areas identified at Homestead Bay, Otago.

Wet Area	Plot Number	Diet Number	Hydrophytic Vegetation Test	Hydric Soils	Hydrology	RMA Wetland	NPS-FM Natural		
Number		Rapid Test	Dominance Test	Prevalence Index	Hydric Vegetation	nyaric soils	nyurology	RIVIA Wetland	Inland Wetland
	Plot 1	Fail	Pass (67%)	Pass (2.67)	Yes	Yes	Yes	Yes	Yes
Wet Area 1	Plot 2	Fail	Uncertain (50%)	Fail (3.91)	No	No	No	No	No
	Plot 3	Fail	Pass (100%)	Pass (2.20)	Yes	Not assessed	Not assessed	Yes	Yes
	Plot 4	Pass	Pass (100%)	Pass (2.0)	Yes	Not assessed	Not assessed	Yes	Yes
	Plot 5	Fail	Fail (33%)	Fail (3.13)	No	Yes	No	No	No
Wet Area 2	Plot 6	Fail	Uncertain (50%)	Fail (3.36)	No	Not assessed	Not assessed	No	No
	Plot 7	Fail	Fail (33%)	Fail (3.07)	Uncertain	Uncertain	No	No	No
	Plot 8	Fail	Fail (0%)	Fail (3.99)	No	Not assessed	Not assessed	No	No
Wet Area 3	Plot 9	Fail	Uncertain (50%)	Fail (3.86)	Uncertain	Yes	Yes	Yes	Yes
Wet Area 4	Plot 10	Fail	Uncertain (50%)	Pass (2.26)	Yes	Uncertain	Yes	Yes	Yes
wet Area 4	Plot 11	Fail	Fail (0%)	Fail (4.15)	No	Not assessed	Not assessed	No	No
	Plot 12	Fail	Fail (0%)	Fail (4.09)	No	Not assessed	Not assessed	No	No
Wet Area 5	Plot 13	Fail	Fail (0%)	Fail (4.91)	No	No	No	No	No
Wet Area 6	Plot 14	Fail	Not assessed (no vegetation present)	Not assessed (no vegetation present)	No	Uncertain	Yes	Yes	Yes
	Plot 15	Fail	Fail (0%)	Fail (4.54)	No	Not assessed	Not assessed	No	No
Wet Area 7	No plot assessment	Pass	-	-	Not assessed	Not assessed	Not assessed	Yes	No
Wet Area 8	No plot assessment	-	-	-	Not assessed	Not assessed	Not assessed	No	No
Wet Area 9	Plot 16	Fail	Pass (100%)	Fail (3.01)	Uncertain	Uncertain	Yes	Yes	Yes



#### 5.2 Wet Area 1

Wet Area 1 is at the northern end of the property and consists of mainly exotic pasture grassland and features a pond, a very small gully, rushlands and an area of pugged bare soils.

#### Plot 1

The rapid test included Yorkshire fog (*Holcus lanatus*; FAC), browntop (*Agrostis capillaris*; FACU), soft rush (*Juncus effusus*; FACW) and sharp spike sedge (*Eleocharis acuta*; OBL) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. Plot 1 was placed within the soft rush and sharp spike sedge rushland (Plate A3-1). The vegetation assessment within this plot, resulted in hydrophytic vegetation being present (Dominance Test = 67%; Prevalence Index = 2.67). Both hydrophytic vegetation tests passed and technically no further wetland tests such as soils and hydrology are necessary. As this, was the first wetland assessment on the property soils and hydrology tests were performed to gain a better understanding of the soils in the area. As expected, these assessments recorded hydric soils and wetland hydrology features being present.

This plot is within a soft rushland that occurs on a slight slope above a pond (Plate A3-16). It is unknown if the pond has been artificially made. It may have historically been a wetland but after many years of farming is now consistently a pond. The pond is just visible in a Retolens image on the 22 April 1964 (SN1641). The pond is consistently inundated, visible in Google Earth from 1 August 2006 to present day. If the pond has been artificially made then the rushland is not a 'natural inland wetland' as it has developed around a deliberately constructed water body (Clause 'c'. within the RMA definition of a natural inland wetland). However, there is no found evidence that this pond has been artificially made and the pond was likely to have been a wetland historically. Therefore, this rush is a natural inland wetland.

#### Plot 2

The rapid test included Yorkshire fog (FAC), Californian thistle (*Cirsium arvense*; FACU), shepherds purse (*Capsella bursa-pastoralis*; no wetland status, so updated to UPL) and sweet vernal (*Anthoxanthum odoratum*; FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. This area contained two small areas of concave, mostly unvegetated soil that had some pugging within it (Plate A3-2). The vegetation assessment for this plot resulted in the Dominance Test (50%) and Prevalence Test (3.91) failing. Due to the marginal Dominance Test result, more assessments were undertaken The soil profile had mineral soils and failed the hydric soils test. There were no hydrological indicators present. This area is not a wetland.

#### Plot 3

The rapid test included Yorkshire fog (FAC), soft rush (FACW), and floating sweetgrass (*Glyceria fluitans*; OBL) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. This plot was placed in a wet channel containing mostly soft rush, Yorkshire fog (*Holcus lanatus*) and floating sweetgrass (*Glyceria declinata*) (Plate A3-3). The vegetation assessment for this plot passed the vegetation test (Dominance Test = 100%; Prevalence Index = 2.20), and contains hydrophytic vegetation. No further testing was necessary and this area is considered a natural inland wetland.



#### 5.3 Wet Area 2

Wet Area 2 is near the southern end of the property and lays just above one of the gullies. This area consists of exotic pasture grassland, rushlands and a concave oval hollow.

#### Plot 4

The rapid test included kneed foxtail (*Alopercurus geniculatus*; FACW) and *Lobelia perpusilla* (FACW) as the dominant species and resulted in a 'passed' result. According to the Wetland Delineation Protocols, no more assessments are required and this area can be considered as a natural inland wetland. However, more hydrophytic vegetation tests were undertaken to ensure a thorough assessment was performed. Plot 4 was placed in a concave oval hollow with abundant kneed foxtail and patches of *Lobelia perpusilla* (Plate A3-4). This plot resulted in a hydrophytic vegetation being present by passing both vegetation tests (Dominance Test = 100%, Prevalence Index = 2.0), therefore this area is a wetland. No further soil and hydrology assessments were required for this plot. This wet area is a natural inland wetland.

#### Plot 5

South of the concave hollow, is a leafless/wiwi rush (*Juncus australis*) rushland. The rapid test included wi/leafless rush (*Juncus australis*; FACW), browntop (FACU), and ryegrass (*Lolium perenne*; FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. Plot 5 was placed within the rushland (Plate A3-5). Vegetation assessments resulted in a fail (Dominance Test = 33%, Prevalence Test = 3.13). However, because the prevalence test was marginal, a soil and hydrology assessment were also undertaken. The soil assessment resulted in hydric soils and the hydrology assessment failed. According to the Wetland Delineation Protocols (Ministry for the Environment 2022), this would mean that this is a drained wetland or atypical environment and further analysis is required. After re-reviewing Google Earth historic imagery and marginal assessment outcomes, it is concluded that this area is not a natural inland wetland.

#### Plot 6

North of the concave oval hollow, is a very small soft rushland on a slope (Plate A3-6). Within the rushland is a shallow drain. The rapid test included soft rush (FACW), white clover (*Trifolium repens*; FACU) and browntop (FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. This plot failed both of the vegetation tests (Dominance Test = 50%, Prevalence Index = 3.36), and is therefore not a wetland.

#### Plot 7

Another plot was undertaken within the same vegetation area of Plot 5 to try and give more clarity to this area (Plate A2-7). The rapid test included wiwi/leafless rush (FACW), white clover (FACU) and browntop (FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. The vegetation assessments results were similar to Plot 5 and resulted in a marginal fail for the Prevalence Index (Dominance Test = 33%, Prevalence Index = 3.07). The soil assessment resulted in uncertain hydric soils and the hydrology assessment failed. It is concluded that this area is not a wetland.



#### 5.4 Wet Area 3

Wet Area 3 is north east of Wet Area 2 and covers an area of exotic pasture grassland and minor undulating land.

#### Plot 8

The rapid test included ryegrass (FACU) and white clover (FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. Plot 8 was placed in a flat area in a slight depression that is covered in exotic herbs and grasses (Plate A3-8). The vegetation tests both failed for this plot (Dominance Test = 0%, Prevalence Index = 3.99). No further assessments were done for this plot and is concluded that it is a dryland habitat and not within a wetland.

#### Plot 9

A deeper oval depression is present just northeast of the above area (Plate A3-2). The rapid test included swamp plantain (*Plantago australis*; FAC), nettle (*Urtica urens*; no wetland status, updated to UPL) and fathan (*Chenopodium album*; no wetland status, updated to UPL) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. The vegetation assessment for this plot resulted in uncertain results, with a Dominance Test of 50% and a Prevalence Index of 3.86. More tests (soil and hydrology) were undertaken to further investigate the wetland assessment. The soil profile presented several signs of hydric soil characteristics. The hydrology assessment also contained several indicators such as being sparsely vegetated concave surface (2H; primary indicator), saturation in aerial imagery (3F; secondary indicator) and geomorphic position (4B; secondary indicator). Plot 9 is within a wetland and can be also classed as a natural inland wetland.

#### 5.5 Wet Area 4

Wet Area 4 is at the southern end of the property and occurs just above one of the larger gullies. This area consists of exotic pasture grassland, rushlands, a concave oval hollow and undulating land.

#### Plot 10

Another deep oval depression is present at the southern end of the property (Plate A3-10). The rapid test included floating sweetgrass (OBL), browntop (FACU) and clammy goosefoot (*Dysphamia pumilio*; no wetland status, updated to UPL) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. The vegetation assessment resulted in uncertain results, with a Dominance Test of 50% (fail) and a Prevalence Index of 2.26 (pass). More tests (soil and hydrology) were undertaken to further investigate the wetland assessment. The soil profile presented several signs of hydric soil characteristics. The hydrology assessment had similar results as Plot 9 and also contained several indicators to confirm wetland hydrology being present. Plot 10 is within a wetland and can be classed as a natural inland wetland.

#### Plot 11

Adjacent to the oval depression above is a small flat area with minimal vegetation (Plate A2-11). The rapid test included Scotch thistle (*Cirsium vulgare*; FACU), ryegrass (FACU), and annual poa (*Poa annua*; FACU) and fathen (no wetland status, updated to UPL) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. This plot failed both of the vegetation tests (Dominance Test = 0%, Prevalence Index = 4.15) and is a dryland area and not a natural inland wetland.



#### Plot 12

This plot is within a small undulation to the north of the large deep oval depression (Plot 10) (Plate A2-12). The rapid test included Californian thistle (FACU) and ryegrass (FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. This plot failed both vegetation tests (Dominance Test = 0%, Prevalence Index = 4.09) and is a dryland area and not a wetland.

#### 5.6 Wet Area 5

Wet Area 5 is a small area that is within the start of the small gully. The area consists of exotic pasture grassland and sweet briar (*Rosa rubiginosa*).

#### Plot 13

This plot is at the bottom of a small mostly unvegetated gully (Plate A3-13). The rapid test included shepherd's purse (no wetland status, updated to UPL), nettle (no wetland status, updated to UPL) and ryegrass (FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. This plot failed both of the vegetation tests (Dominance Test = 0%, Prevalence Index = 4.91) and is a dryland area and not a natural inland wetland.

#### 5.7 Wet Area 6

This potential wetland is a flat to shallow mostly unvegetated oval depression. Only a small area had pugging and surface soils cracks nearby the fenceline. The rest of the area had compacted soils and sparse exotic herbs and grasses around the edges. Unvegetated areas within paddocks can also be caused from having more intense congregation from stock. However, since the area was in a depression and showed some hydrological indicator signs, the area was assessed.

#### Plot 14

The rapid test included fathen (no wetland status, updated to UPL), dwarf mallow (*Malva neglecta*; no wetland status, updated to UPL), white clover (FACU) and ryegrass (FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. Plot 14 was placed within the small pugged area of the lowest part of the mild depression (Plate A3-14). This area was a difficult assessment as no plant species were within the plot, meaning that the vegetation assessment could not be undertaken. The soil profile contains some low chroma colours and iron concretions that are suggestive of potential hydric soils. The topsoil chroma is 3/2, and any topsoils 3 or less are not good indicators of hydric soils as many topsoils have this colour range (Fraser *et al.* 2018). This soil profile has an outcome of 'uncertain' soils. The hydrology assessment contained indicators such as being sparsely vegetated concave surface (2H; primary indicator), saturation in aerial imagery (3F; secondary indicator) and geomorphic position (4B; secondary indicator). According to the Wetland Delineation Protocols (Ministry for the Environment), if the soil assessment fails (uncertain here), and the hydrology passes (this area contains one primary indicator and also two secondary hydrology indicators), then this area is a wetland. This area is likely to be a very recent natural inland wetland.

#### Plot 15

The rapid test included fathen (no wetland status, updated to UPL), dwarf mallow (no wetland status, updated to UPL), white clover (FACU) and ryegrass (FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were



required. A plot was placed on the upper edge of the depression where the soil was compacted and contained a few sparse exotic herbs and shrubs (Plate A3-15). Both vegetation assessments failed for this plot (Dominance Test = 0%, Prevalence Index = 4.54). This area is not a natural inland wetland.

#### 5.8 Wet Area 7

Wet Area 7 is a large area that has possibly been dug out and contains an island in the middle (Plate A3-17 and A3-18). This area may have historically been a wetland but was modified by the landowner. Google Earth images from 1 August 2006, suggests that the modification may have occurred around this date.

The large hole has steep sides with exotic pasture species. The bottom on the hole contains a herbfield of marsh bedstraw (*Galium palustre* subsp. *palustre*; OBL), sharp spike sedge (OBL), kneed foxtail (FACW), waoriki (*Ranunculus glabrifolius*; OBL) and Shepard's purse (no wetland status). The dominate species present (marsh bedstraw, sharp spike sedge, kneed foxtail and waoriki) all have a wetland status of either FACW and OBL. This area passes the rapid test and is considered a wetland under the RMA. This wetland is excluded from the 'natural wetland' definition because it has been formed in a constructed excavation and was likely intended to be a pond (currently dry).

#### 5.9 Wet Area 8

This area covers undulating land covered in exotic pasture grassland and a terrace. One area at a low point within the undulating land looked to may have contained water-logging in the initial and brief look at historic Google Earth images. This area contains consistent, well-covered exotic pasture species that are mostly Facultative Upland or Upland species (dryland species).

The other area considered to be worth checking out after the initial and brief look at historic Google Earth images was a low-lying area at the bottom of the shrubland terrace. This area contained species that are all Facultative Upland species such as porcupine shrub (*Melicytus alpinus*), barley grass (*Hordeum murinum* subsp. *murinum*), cocksfoot (*Dactylus glomerata*) and Scotch thistle (*Cirsium vulgare*). The other species present such as burdock (*Arctium lappa*), dovesfoot cranesbill (*Geranium molle*), nettle and dwarf mallow do not have a wetland status but all are likely to be Facultative Upland or Upland species (dryland species). The dominance of dryland vegetation species and the lack of any hydrology features leads to the conclusion that this area is a dryland.

#### 5.10 Wet Area 9

Wet Area 9 is within an ephemeral stream just north of the airstrip. The ephemeral stream has eroded a wide channel on the landscape and has steep banks containing bare soils, rocks and exotic pasture grasses. The flatter stream bed, has signs of pugging damage and is mostly covered with exotic pasture grasses, featuring some deeper incised gravel channels.

#### Plot 16

The rapid test included Yorkshire fog (FAC), ryegrass (FACU), white clover (FACU) and browntop (FACU) as the dominant species and resulted in a 'failed' result. Due to the failed result of the rapid test more hydrophytic vegetation assessments were required. Plot 16 was placed within the small pugged area of the ephemeral stream bed (Plate A3-19).

The vegetation assessment resulted in uncertain results, with a Dominance Test of 100% (pass) and a Prevalence Index of 3.01 (fail). More tests (soil and hydrology) were undertaken to further investigate the wetland assessment. The soil profile resulted with an uncertain outcome. The hydrology assessment had one primary hydrology indicator present (oxidised rhizophere on roots), to confirm



wetland hydrology being present. Plot 16 is within a wetland and can be classed as a natural inland wetland.

### 6.0 Natural inland wetlands

#### 6.1 Overview

Seven natural inland wetlands have been identified at the Homestead Bay property proposed for development (Figure 2). The natural inland wetlands are small in size and are predominantly dominated by exotic plant species. Three classes of natural inland wetlands (marsh, swamp and ephemeral) are present at the Homestead Bay property. Despite, the exotic plant dominance the wetlands are still valuable in their ability to be a carbon sink and providing habitat for indigenous bird and invertebrates species, in particular wading birds such as pūkeko (*Porphyrio melanotus;* Not Threatened).

Four of natural inland wetlands present are ephemeral wetlands. Ephemeral wetlands are typically in closed depressions and contain low statue plant species that are often arranged in a zonation pattern. This type of wetland has unique hydrology characteristics by being intermittently inundated throughout the year. Typical hydrology of an ephemeral wetland is usually ponded during winter/spring, with the water level gradually lowering in summer (Johnson and Rogers 2003). Ephemeral wetlands are ecologically valuable as they can contain a diverse range of plant species, host a high proportion of uncommon and threatened plants (Johnson and Rogers 2003) and provide habitat for wading indigenous birds. Ephemeral wetlands threatened by many factors including human-induced modifications, sedimentation, nutrient enrichment, pollutants, trampling impacts from mammals, weed invasions and introduced fish (Johnson and Rogers 2003). Ephemeral wetlands are a critically endangered naturally uncommon ecosystem in New Zealand (Holdaway *et al.* 2012)

#### Soft rush-browntop-sharp spike sedge rushland marsh (0.0419 hectares)

Clusters of soft rush are spread throughout a mosaic of browntop, sharp spike sedge and Yorkshire fog (Plate 1). Other species occasionally present include white clover, wīwī/leafless rush and jointed rush (*Juncus articulatus*). A sedge species is also present in very low abundance. The sedge did not have an inflorescence to confirm identification but is likely Sinclair's sedge (*Carex sinclairii*), due to its leaf characteristics<sup>1</sup> and habitat match.

٠

<sup>&</sup>lt;sup>1</sup> Double folded leaf, leaf width of 4-4.5 millimetres, leaf length of c. 40 centimetres and light yellow-green leaf colouration.





Plate 1 – Rushland marsh wetland above the pond in Wet Area 1 at Homestead Bay. 30 January 2025.

#### 2. Yorkshire fog marsh

Within one of the ephemeral streams is a small marsh wetland. The marsh wetland is dominated by Yorkshire fog and has the other occasional exotic herb and grass within it. It is surrounded by drier areas dominated with browntop.

#### 3. Soft rush-floating sweet grass-Yorkshire fog grassland swamp (0.0087 hectares)

This swamp wetland lays in a channel that sits below the pond in Wet Area 1 (Plate 2). The swamp wetland contains a mosaic of soft rush, Yorkshire fog, floating sweetgrass, curled dock (Rumex crispus) and jointed rush. The wetland contains the occasional clusters of water fern (Azolla rubra) sitting on the water surface.

#### 4. Kneed foxtail-Lobelia perpusilla grassland ephemeral wetland (0.0372 hectares)

This ephemeral wetland is located in the middle of the Wet Area 2 and is a concave hollow that is lined with soft rush on the northern end and leafless/wiwi rush on the southern end (Plate 3). The ephemeral wetland contains abundant kneed foxtail with common patches of the indigenous herb Lobelia perpusilla.









Plate 2 – Rushland swamp wetland below the pond in Wet Area 1 at Homestead Bay. Plot 3 can be seen within the wetland. 30 January 2025.



Plate 3 – Ephemeral wetland surrounded by rushlands in the middle of Wet Area 2 at Homestead Bay. 30 January 2025.



#### 5. [Swamp plantain] herbfield ephemeral wetland (0.0130 hectares)

Another ephemeral wetland is located in Wet Area 3. This wetland is within an oval depression and is mostly unvegetated (Plate 4). A few exotic species are scarce and scattered near the edges and include: marsh plantain, ryegrass, nettle and clammy goosefoot.



Plate 4 – Ephemeral wetland within the exotic pasture grassland within Wet Area 3 at Homestead Bay. 30 January 2025.

## 6. [Floating sweetgrass-kneed foxtail-clammy goosefoot] grassland ephemeral wetland (0.0289 hectares)

This ephemeral wetland is also within a mostly unvegetated oval depression (Plate 5). Scarce and scattered exotic species located near the edges of the wetland include floating sweetgrass, browntop, clammy goosefoot, white clover, and black nightshade (*Solanum nigrum*).

#### 7. Mudflat ephemeral wetland (0.0105 hectares)

This unvegetated ephemeral wetland is lies in a small depression and is very small in size. This area is likely a recently developed wetland, as suggested by the 'uncertain' soil result in the guidelines (Fraser *et al.* 2018). The first evidence of water pooling in this wetland on Google Earth can only be seen from 10 February 2010. The ephemeral wetland is at the lowest point (paler soil) of a compacted open soil patch (Plate 6).





Plate 5 – Ephemeral wetland within the exotic pasture grassland within Wet Area 4 at Homestead Bay. 3 February 2025.



Plate 6 – Ephemeral wetland within the exotic pasture grassland within Wet Area 6 at Homestead Bay. 4 February 2025.



### 7.0 Summary

Nine areas were investigated for the possibility of wetlands being present. A total of seven natural inland wetlands were identified within the property at Homestead Bay proposed for a sub-division development. The marsh and swamp wetlands are dominated by exotic species, with soft rush being the most common species present. The ephemeral wetlands are mostly dominated by exotic plant species, nonetheless still contain typical characteristics of their ecosystem type. These natural inland wetlands will likely provide habitat to indigenous bird and invertebrate species. The ephemeral wetlands are particularly ecologically valuable due their potential to host diverse plant species as well as uncommon and threatened plant species. Ephemeral wetlands are a unique and uncommon habitat and has been listed as a critically endangered ecosystem. All wetland present are important on a national scale as New Zealand has lost 90% of its pre-human wetland extent, making wetlands the most nationally threatened ecosystem type (Aussiel *et al.* 2008).

The residential development could consider avoiding these wetlands and incorporating them into their reserve areas. All wetlands would benefit from enhancement actions such as indigenous plantings (around the wetland margin for the ephemeral wetlands), weed control and cattle exclusion. Light grazing from sheep can be beneficial for ephemeral wetlands as they discourage woody weed species and encourage low-statue turf communities that are typical for this type of wetland ecosystem.

## Acknowledgments

Amanda Leith (Planner/Director of Remarkables Planning) is thanked for providing useful information and site access. Farm Managers Dave Atkin and Matt Little are thanked for allowing access to the active farm areas.

## References

Atkinson I.E. (1985): Derivation of vegetation mapping units for an ecological survey of Tongariro National Park, North Island, New Zealand. New Zealand Journal of Botany 23: 361-378.

Ausseil A-G., Gerbeaux P., Chadderton L., Stephens T., Brown D. and Leathwick J. (2008): *Wetland ecosystems of national importance for biodiversity: Criteria, methods and candidate list of nationally important inland wetlands*. Prepared for Department of Conservation. Landcare Research Contract Report: LC0708/158. 172 pp

Clarkson R.C, Fitzgerald N.B, Champion P.D., Forester L., and Rance B.D. (2021): *New Zealand Wetland Plant List 2021*. This report was prepared by Manaaki Whenua – Landcare Research for Hawke's Bay Regional Council. 66 pp.

Clarkson B.R. (2013): A vegetation tool for wetland delineation in New Zealand. Landcare Research, Hamilton. 70 pp.

Clarkson B.R., Sorrell B.K., Reeves P.N., Champion P.D. Partridge T.R. and Clarkson B.D. (2003): *Handbook for monitoring wetland condition* (Revised October 2004). Coordinated Monitoring of New Zealand Wetlands. A Ministry for the Environment Sustainable Management Fund Project (5105). 73 pp.

Cosgrove G, Dodd M, James T. (2022): *National list of exotic pasture species*. Prepared for the Ministry for the Environment by AgResearch Ltd. Wellington: Ministry for the Environment. 24 pp.



de Lange, P.J., Gosden, J., Courtney, S.P., Fergus, A.J., Barkla, J.W., Beadel, S.M., Champion, P.D., Hindmarsh-Walls, R., Makan, T. and Michel, P. (2024): *Conservation status of vascular plants in Aotearoa New Zealand, 2023*. New Zealand Threat Classification Series 43. Department of Conservation, Wellington. 105 pp.

Fraser S., Singleton P., and Clarkson B. (2018): *Hydric soils – field identification guide*. Manaaki Whenua – Landcare Research contract report prepared for Tasman District Council. 75 pp.

Holdaway R.J., Wiser S.K. and Williams P.A. (2012): *Status Assessment of New Zealand's Naturally Uncommon Ecosystems*. Society for Conservation Biology. Conservation Biology, Volume 26, No. 4, 619-629. 10 pp.

Johnson P.N. and Gerbeaux P. (2004): *Wetland types in New Zealand*. Department of Conservation, Wellington. 184 pp.

Johnson P. and Rogers G. (2003): Ephemeral wetlands and their turfs in New Zealand. Science for Conservation 230. Department of Conservation, Wellington. 88pp.

McEwen W.M. (1987): *Ecological Regions and Districts of New Zealand*. Third revised edition in four 1:500,000 maps. Publication No. 5 (in four parts). Part 3. New Zealand Biological Resources Centre, Department of Conservation, Wellington. 137 pp.

MetService New Zealand (2025): Monthly rainfall and historical average rainfall for Queenstown airport.

Retrieved from: <a href="https://www.metservice.com/towns-cities/regions/southern-lakes/locations/queenstown/past-weather">https://www.metservice.com/towns-cities/regions/southern-lakes/locations/queenstown/past-weather</a>, February 2025.

Ministry for the Environment (2020): *Wetland delineation protocols.* Wellington: Ministry for the Environment. 14 pp.

Ministry for the Environment (2020): *National Policy Statement for Freshwater Management 2020*. Wellington: Ministry for the Environment. 70 pp.

Ministry for the Environment (2020): *Resource Management (National Standards for Freshwater) Regulations*. Wellington: Ministry for the Environment. 62 pp.

Ministry for the Environment (2021): *Wetland delineation hydrology tool for Aotearoa New Zealand*. Wellington: Ministry for the Environment. 83 pp.

Ministry for the Environment (2021): Defining 'natural wetlands' and 'natural inland wetlands'. Wellington: Ministry for the Environment. 25 pp.



## Plant species recorded and their wetland indicator status<sup>2</sup>

Species	Common Name	Threat Status <sup>3</sup>	Status	Wetland Indicator Status
Achillea millefolium	Yarrow	-	Exotic	FACU
Agrostis capillaris	Browntop	-	Exotic	FACU
Alopecurus geniculatus	Kneed foxtail	-	Exotic	FACW
Anthoxanthum odoratum	Sweet vernal	-	Exotic	FACU
Arctium lappa	Burdock	-	Exotic	None
Azolla rubra	Pacific azolla, azolla, red	Not	Indigenous Non-	None
	azolla	Threatened	Endemic	
Capsella bursa-pastoris	Shepherds purse		Exotic	None
Carex sp. <sup>4</sup>	Sedge	Not Threatened	Indigenous Non- Endemic	None
Carex sp.	Sedge	-	Indigenous Non- Endemic	None
Cerastium glomeratum	Chickweed	-	Exotic	FACU
Chenopodium album	Fathen	-	Exotic	None
Cirsium arvense	Californian thistle	-	Exotic	FACU
Cirsium vulgare	Scotch thistle	-	Exotic	FACU
Cynosurus cristatus	Crested dogstail	-	Exotic	UPL
Dactylis glomerata	Cocksfoot	-	Exotic	FACU
Dysphania pumilio	Clammy goosefoot	-	Exotic	None
Eleocharis acuta	Spike sedge	Not Threatened	Indigenous Non- Endemic	OBL
Erodium cicutarium	Storksbill	-	Exotic	None
Galium palustre	Marsh bedstraw	-	Exotic	OBL
Geranium molle	Dovesfoot cranesbill	-	Exotic	None
Glyceria fluitans	Floating sweetgrass	-	Exotic	OBL
Holcus lanatus	Yorkshire fog	-	Exotic	FAC
Hordeum murinum	Barley grass	-	Exotic	FACU
Hypericum humifusum	Trailing Saint John's wort	-	Exotic	FAC
Hypochaeris radicata	Catsear	-	Exotic	FACU
Juncus articulatus	Jointed rush	-	Exotic	FACW
Juncus australis	Wīwī, wī, leafless rush	Not Threatened	Indigenous Non- Endemic	FACW
Juncus effusus	Soft rush	-	Exotic	FACW
Juncus tenuis subsp.		-	Exotic	FACW
dichotomus				
Leontodon saxatilis	Hawkbit	-	Exotic	FAC
Lobelia perpusilla		Not Threatened	Indigenous Endemic	FACW
Lolium perenne	Ryegrass	-	Exotic	FACU

<sup>&</sup>lt;sup>2</sup> Any species with a 'none' status was changed to a 'OBL' status in the assessment.

<sup>&</sup>lt;sup>3</sup> de Lange *et al*. 2024

<sup>&</sup>lt;sup>4</sup> Likely *Carex sinclairii* 



Species	Common Name	Threat Status <sup>3</sup>	Status	Wetland Indicator Status
Malva neglecta	Dwarf mallow	-	Exotic	None
Melicytus alpinus	Porcupine shrub	Not Threatened	Indigenous Endemic	FACU
Plantago australis	Swamp plantain	-	Exotic	FAC
Poa annua	Annual poa	-	Exotic	FACU
Ranunculus glabrifolius	Waoriki	Not Threatened	Indigenous Non- Endemic	OBL
Rumex crispus	Curled dock	-	Exotic	FAC
Rumex obtusifolius	Broad-leaved dock	-	Exotic	FAC
Sagina procumbens	Pearlwort	-	Exotic	FACU
Solanum nigrum	Black nightshade	-	Exotic	FACU
Sonchus asper	Prickly puha	-	Exotic	FACU
Spergula arvensis	Spurrey	-	Exotic	None
Taraxacum officinale	Dandelion	-	Exotic	FACU
Trifolium repens	White clover	-	Exotic	FACU
Urtica urens	Nettle	-	Exotic	None
Verbascum thapsus	Woolly mullein	-	Exotic	None
Verbascum virgatum	Moth mullein	-	Exotic	None



### Monthly rainfall data for 2024, 2025 and historical averages





### Site photographs



Plate A3-1 – Vegetation Plot 1 within Wet Area 1 at Homestead Bay. 30 January 2025.



Plate A3-2 – Vegetation Plot 2 within Wet Area 1 at Homestead Bay. 30 January 2025.



Plate A3-3 – Vegetation Plot 3 within Wet Area 1 at Homestead Bay. 30 January 2025.



Plate A3-4 – Vegetation Plot 4 within Wet Area 2 at Homestead Bay. 30 January 2025.





Plate A3-5 – Vegetation Plot 5 within Wet Area 2 at Homestead Bay. 30 January 2025.



Plate A3-6 - Vegetation Plot 6 within Wet Area 2 at Homestead Bay. 30 January 2025.



Plate A3-7 – Vegetation Plot 7 within Wet Area 3 at Homestead Bay. 3 February 2025.



Plate A3-8 - Vegetation Plot 8 within Wet Area 3 at Homestead Bay. 3 February 2025.



Plate A3-9 – Vegetation Plot 9 within Wet Area 3 at Homestead Bay. 3 February 2025.



Plate A3-10 – Vegetation Plot 10 within Wet Area 4 at Homestead Bay. 3 February 2025.





Plate A3-11 – Vegetation Plot 11 within Wet Area 4 at Homestead Bay. 3 February 2025.



Plate A3-12 –Vegetation Plot 12 within Wet Area 4 at Homestead Bay. 3 February 2025.



Plate A3-13 – Vegetation Plot 13 within Wet Area 5 at Plate A3-14 – Vegetation Plot 14 within Wet Area 6 at Homestead Bay. 4 February 2025.



Homestead Bay. 4 February 2025.



Homestead Bay. 4 February 2025.



Plate A3-15 – Vegetation Plot 15 within Wet Area 6 at Plate A3-16 – Pond in Wet Area 1 at Homestead Bay. 30 January 2025.





Plate A3-17 – Drain on the upper edge of the large excavation area within Wet Area 7 at Homestead Bay. Area 7 at Homestead Bay. 30 January 2025. 30 January 2025.

Plate A3-18 – The large excavation area within Wet



Plate A3-19 - Plot 16 within Area 9 at Homestead Bay. 3 September 2025.



## Wetland delineation plot sheets

NEW ZEAL	AND WETLA	ND DELINEAT	ION DATA FORM
S	ECTION A -	SITE INFORMA	ATION
Owner: RCL group  Landform: Undulating land  Is the land drained (circle) YES (NO)*  GPS (NZTM): F126 2952 N4998  *Small pond nearby (possible)  Are climatic/hydrologic conditions on the site type  Are vegetation, soil or hydrology significantly dist  Are vegetation, soil or hydrology naturally proble  SUMMARY OF FINDINGS—Attach site  Hydrophytic vegetation present? YES  Hydric soils present?  Wetland hydrology present?  YES	Local relief: Vory Investigator(s): FA  894 Alt  7 mm - made) ical for this time of year urbed? (circle) matic? (circle) map showing samp	in Gallagher itude m: 388 m  Mis may be  ? YES NO (circle ap Are 'normal circumsta Explain answers in Re	Land use: <u>ullivated</u> & grazed  Land cover; <u>lxotic</u> <u>grass/and</u> Soil*C: <u>Not rewelled</u> . Slope*: <u>mild grad</u> Photo Nos: <u>sared</u> to <u>falder</u> drawing the area.  propriate; if NO explain in Remarks)  unces' present? (circle) YES NO  marks if needed  ransects, important features etc.
wettand nydrology present: 123 1		B – VEGETATIO	ON
Use scientific names of plants.	ute Dominant ver Species?	Indicator laylure \$\frac{1}{5}\$ Status species? \$\frac{1}{5}\$	Dominance Test:
1. Holcus lanatus 25 2. Agrotis capillaris 40 3. ELE aca 30 4. JUNeff 10 5. JUN art 3 6. TRIrep 0 7. Carex sp> CARsin 2 8. 9. 10. 11. 12. 207=27.1 Total cover = 10. 507=55.25	<u>y</u> <u>y</u> <u>y</u> <u>5                          </u>	FAC Y 25 FACU Y 40 OBL FACW FACW FACW FACU Y 0.5 OBL	Dominance Test is >50%  Prevalence Index is ≤3.0¹  Morphological adaptations¹ (supporting data in Remarks)  Problematic hydrophytic vegetation¹  ¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic  Hydrophytic vegetation present?  YES  NO  UNCERTAIN  UNCERTAIN
Remarks: PITV(= 59% exotic Carex sp. is highly which makes iden match that of CA	likely to be hitication do	Carex sinclaini	. There was no influrescence haracheristics and habitat



	Matrix colour (moist)	Mottles colour (moist)	Mottles %1	Mottles Size <sup>2</sup>	Mottle location <sup>3</sup>	Material <sup>4</sup>	Remarks
0-28	4/1	4/3	5	very line	matrix	mineral	
28 - 38	4/3	4/1	20	Medium	matrix	mineral	
•							
	charts; 'Use size class I indicators:	es; 'Ped face, pore, with Soil drainage (circle)				ty), humic, mineral soi	
Organic laye	ers: (	Concretions:		Colours: profile	In	cation: Depression Fla	
=/	soil material	Iron concretions		Gley OR		eter table: Depth (cm)	
Litter		Manganese concre		Mottled		sh GW (Perched) Seep ns: Depth (cm )	age Tual Little
Mesic	L	Consistence:		Reductimo	D-		- si- Duri- Fragi Ortstein
Humic		Plastic		Redox mot	led Lay	vers: Depth (cm)	
Peaty to		Sticky Fluid		Redox segre	-gations	w perm argillic Pugged	
Hydric soil	s present?	YES I	NO $\square$		ERTAIN	NZSC subgroup	
Secondary Water- Drainag	posits (2C)  / hydrology indic  stained leaves (2K)  ge patterns (2L)  ason water table (3E)  tion in aerial imagery	SI	of <u>2</u> require eomorphic p hallow aquita AC-neutral te	osition (4B) ard (4C)	boxes that apply  FAC-neutral test  1. No. OBL & FA	(4D); refer to Section E CW dominant species PL dominant species	
Skatch of site	ydrology present			NO			
	unullini.	Metal College	an May	MA M.	Muren	pond	June June
- 419							
	lore grassian	d	ruskla	md	parture g	rass land	lot x



#### NEW ZEALAND WETLAND DELINEATION DATA FORM SECTION A - SITE INFORMATION Site: Homestead Bay Region: Dtago Sampling point: WA1 - plot 2 Owner: RCL group Date: 30/1/2025 \_ Land use: altirated & grazed Landform: undulating land Local relief: concare to flut Land cover: exotic grassland Investigator(s): Frin Gallagher Is the land drained (circle) YES NO Soil °C: not recorded Slope :: concare GPS (NZTM): E1265993 N 4998871 Altitude m: 389 Photo Nos: saved in report folder Are climatic/hydrologic conditions on the site typical for this time of year? YES NO (circle appropriate; if NO explain in Remarks) Are vegetation, soil or hydrology significantly disturbed? (circle) Are 'normal circumstances' present? (circle) (YES) NO Are vegetation, soil or hydrology naturally problematic? (circle) Explain answers in Remarks if needed SUMMARY OF FINDINGS—Attach site map showing sampling point locations, transects, important features etc. Hydrophytic vegetation present? YES NO Is the sampled area within a wetland? YES Hydric soils present? YES NO Wetland hydrology present? YES NO SECTION B – VEGETATION Use scientific names of plants. Absolute Dominant Dominance Test: Tree Stratum (Plot size:\_\_\_\_ % cover Species? No. Dominant Spp. OBL/FACW/FAC (A) Tot. Dominant Spp. across strata (B) \_\_\_\_\_\_\_ % OBL/FACW/FAC (A/B) 50 Prevalence Index: Total % cover of: Multiply by: Total cover = OBL O x1=\_0 Sapling/Shrub Stratum (Plot size:\_\_\_\_ FACW \_\_0 x2=\_\_0 x3=\_9 FACU \_ 6.3 x4 = 25.2Total \_\_\_\_\_\_\_\_(A) 44.2 (B) Prevalence Index (B/A) = 3.9/ Total cover = Herb Stratum (Plot size: 2m2) Hydrophytic vegetation indicators: 1. Holcus lanatus FAC Dominance Test is >50% 2. Cirsium vulgaris 4 FACU Prevalence Index is ≤3.01 3. Lolium perrenne 0.1 Morphological adaptations<sup>1</sup> (supporting data in Remarks) 4. Capsella bursa-pastoris 2 None - UPL 5. Anthoxanthum radiation 2 Problematic hydrophytic vegetation<sup>1</sup> FACU 6. Cerastrium glomeratum 0.1 7. Sonchus asper <sup>1</sup>Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic Hydrophytic vegetation present? 10.\_ YES 11.\_ 12.\_\_\_ UNCERTAIN Total cover = \_\_ 507, = 5,65 Remarks: Remarks: -pugging and slight surface cracks - no Eurther tests undertaken



#### NEW ZEALAND WETLAND DELINEATION DATA FORM SECTION A – SITE INFORMATION Site: Homestead Bay Region: Otago Sampling point: Plot 2 Owner: RCL Homestead Bay Ltd Date: 3 September 2025 Land use: Farmland - cattle grazing Local relief: Slight slope Land cover: Exotic grassland Landform: Undulating toeslope Investigator(s): Erin Gallagher Soil °C:Not measured Slope :Not measured Is the land drained (circle) YES NO GPS (NZTM):E1265994 N4998871 Altitude m:\_372m Photo Nos: Photos in report folder 7522 Are climatic/hydrologic conditions on the site typical for this time of year? YES NO (circle appropriate; if NO explain in Remarks) Are vegetation, soil or hydrology significantly disturbed? (circle) Are 'normal circumstances' present? (circle) Are vegetation, soil or hydrology naturally problematic? (circle) Explain answers in Remarks if needed SUMMARY OF FINDINGS—Attach site map showing sampling point locations, transects, important features etc. NO Is the sampled area within a wetland? YES Hydrophytic vegetation present? YES Hydric soils present? YES NO NO Wetland hydrology present? YES SECTION B - VEGETATION Use scientific names of plants. Absolute Dominant Indicator Dominance Test: Tree Stratum (Plot size: % cover Species? Status No. Dominant Spp. OBL/FACW/FAC Tot. Dominant Spp. across strata % OBL/FACW/FAC (A/B) Prevalence Index: Total % cover of: Multiply by: Total cover = OBL Sapling/Shrub Stratum (Plot size:\_\_\_\_ FACW FAC FACU UPL Total (A) Prevalence Index (B/A) = Total cover = Herb Stratum (Plot size: Hydrophytic vegetation indicators: Dominance Test is >50% Prevalence Index is ≤3.01 Morphological adaptations1 (supporting data in Problematic hydrophytic vegetation<sup>1</sup> <sup>1</sup>Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic Hydrophytic vegetation present? YES NO UNCERTAIN Remarks: Vegetation assessment notes on a separate form.



		S	ECTIC	NC-	SOIL A	ND H	DROL	OGY	
rofile descrip	otion: (Describe to	the depth need	led to conf	irm indicat	or presence/ab	sence, 30 c	m default)		
Depth (cm)	Matrix colour (moist)	Mottles (mo	Vi	Mottles %1	Mottles Size <sup>2</sup>	Mottle	location <sup>3</sup>	Material <sup>4</sup>	Remarks
)-36cm	3/1							Mineral	
		-			-		-		
1.00									
line 94 anns a	hade. <sup>2</sup> ttes else els	Specifican	nan wish	is and also	to within	matelu <sup>4</sup> 0	enale fans	ty), humic, mineral s	all'
	indicators:					matrix; C		use of wetness (circle	
		Soil draina	ge (circle)		a construction of		Loc		lat Valley Gully Slope
Organic laye	rs: soil material	Iron con	rretions	ľ	Colours: profile		er:	iter table: Depth (cm	
Utter	Jon Hateria		ese concret	tions [	Gley profil Mottled pr		Hig	h GW Perched See	epage Tidal Lithic
Fibric		Nodular			Horizon:	Offic TOTAL	Par	ns: Depth (cm )	<u></u>
Mesic		Consistence:		Í	Reductimo	rphic	Par	Humus Fe-pan De	ensi- Duri- Fragi Ortstein
Humic		Plastic		Ì	Redox mot	V-01 (18)	Res	stricting layer: Depth	(cm)
Peaty to	psoil	Sticky		Ì	Redox segr	egations	Slo	w perm argillic	
Peaty su	ıbsoil	Fluid		Ì	Perch-gley	features		Pugged	
ydric soil:	s present?	YES	N	0 🗸	UNC	ERTAIN		NZSC subgroup	D
Ground Soil sate Water of Sedime Drift de	water (1A) lwater <30 cm (1B) uration <30 cm (1C marks (2A) ent deposits (2B) eposits (2C)	· =	Iron depo Surface so Inundatio Sparsely v Salt crust	oil cracks (2 n on aerial regetated co (21)	F) imagery (2G) oncave surface	Ē	Hydroger Oxidised Reduced Reduced High wat	nvertebrates (2J) n sulphide odour (3A rhizosphere on roots iron (3C) iron in tilled soil (3D) er table stunted/stre	(3B)
_	hydrology ind	icators: min	imum of	<u>z</u> requir	ea; cneck ai			***	
=	tained leaves (2K)				osition (4B)			(4D); refer to Section	1
	e patterns (2L)			llow aquita				CW dominant specie	
_	son water table (38	Planner.		2-neutral te	SEC. 33.5.	3. To		PL dominant species	(B) (A+B)
Saturati	on in aerial imager	y (3F)	Fro	st-neave h	ummocks (4E)	10000	cai C-neutral (>	50%)	(A/A+B)*100
Vetland h	ydrology presei	nt? YE	s 🔲		NO				
ketch of site	/soil:								
temarks:									



	SECTION A - S	SITE INFORMA	TION
Site: Homestead Bay Owner: RCL group Landform: undulating (a) Is the land drained (circle) YES (NO) GPS (NZTM): £ 1265918	Sampling point: WA1 - plot 3 Land use: <u>cultivated &amp; grazed</u> Land cover: <u>exotic grassland</u> Soil °C: <u>not recorded</u> Slope°: <u>gradual</u> it Photo Nos: <u>saved to report</u> tolder		
Are climatic/hydrologic conditions on the Are vegetation, soil or hydrology significate Are vegetation, soil or hydrology natural	ntly disturbed? (circle) y problematic? (circle)	Are 'normal circumstar	
SUMMARY OF FINDINGS—Attac Hydrophytic vegetation present? Hydric soils present? Wetland hydrology present?	YES NO IS		hin a wetland? YES NO
	SECTION I	3 – VEGETATIO	N
Use scientific names of plants.  Tree Stratum (Plot size:)  1. 2. 3. 4.  Total cover =  Sapling/Shrub Stratum (Plot size:)  1. 2. 3. 4.  Total cover =  Herb Stratum (Plot size:)  1. 2. 3. 4. 5.  Total cover =  Herb Stratum (Plot size:)  2.  Junus landus  2.  Junus effusus  3.  Gylleria declinata  4.  Az Olla rubta  5.  Junus artiulatus  6.  Girsium arvense  7.  Rumox crisp us  8.  9.  10.  11.  12.  207, = 16.42  30 7. = 41,02	30 y 35 y 15 0.1 1 0.5 0.5	Indicator Prime & Status Species? Status Species? Status Species & Status Species & Status Species & Speci	Dominance Test:  No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  (B) 2  % OBL/FACW/FAC  Prevalence Index:  Total % cover of:  Multiply by:  OBL
Remarks: No further as tess Possible man-ma			1 channel.



Frome descrip	otion: (Describe to th	e depth needed to con	firm indicate	or presence/ab	sence, 30 cm default		
Depth (cm)	Matrix colour (moist)	Mottles colour (moist)	Mottles %1	Mottles Size <sup>2</sup>	Mottle location <sup>3</sup>	Material⁴	Remarks
							1
lu or 1	. 200	3n 16 W			. 40		
Hydric soil		Soil drainage (circle)				aty), humic, mineral ause of wetness (cir	
Organic layer	s: C	oncretions:		olours: profile	10		Flat Valley Gully Slope
	soil material	Iron concretions		Gley OR		ater table: Depth (c	m) eepage Tidal Lithic ,
Litter	-	Manganese concre Nodular	_	Mottled lorizon:		ans: Depth (cm )	eepage ridai Littiic ,
Mesic	- 0	onsistence:		Reductimo	rphic Pa	an Humus Fe-pan I	Densi- Duri- Fragi Ortstein
Humic Peaty top	esoil	Plastic Sticky		Redox mot		yers: Depth (cm) ow perm argillic	
Peaty sub		Fluid		Redox segri		Pugged	
Hydric soils	present?	YES N	0	UNC	RTAIN	NZSC subgro	up
Primary h	vdrology indica	tors: minimum of	1 require	d: check all	hoxes that apply		
Soil satur Water ma	deposits (2B)	Inundatio	oil cracks (2F n on aerial in regetated co	nagery (2G)	Oxidised Reduced	en sulphide odour (3 d rhizosphere on roo d iron (3C) d iron in tilled soil (3 ter table stunted/st	ots (3B)
econdary h	ydrology indica	tors: minimum of	2 require	d; check all	boxes that appl	у	
Drainage p	ined leaves (2K) patterns (2L) n water table (3E) n in aerial imagery (3	Sha	omorphic po llow aquitar C-neutral tes st-heave hur	d (4C)	1. No. OBL & F/	t (4D); refer to Secti ACW dominant speci JPL dominant speci >50%)	cies(A)
Vetland hyd	rology present?	YES V		NO			
rature gr							Pond
	rushland				x 111 \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	SE SE MILE	W
	swamp					rushla	nd
marks:	y	Log-floa-	Live .	11		1	



SE	CTION A – SI	TE INFORMA	TION
Owner: RCL Croups  Landform: hide gully  Is the land drained (circle) YES NO  GPS (NZTM): £1265293 N4997	Region: Otrogo Date: 30/01/20 Local relief: concar Investigator(s): Emr	ort or Gallagher	Sampling point: WA2 - plot 4  Land use: Cultivated & grazed  Land cover: crotic grassland  Soil *C: not recorded Slope*: n/a  Photo Nos: sared in upont folder
Are climatic/hydrologic conditions on the site typical Are vegetation, soil or hydrology significantly distured are vegetation, soil or hydrology naturally problem SUMMARY OF FINDINGS—Attach site method by the hydrophytic vegetation present? YES Hydric soils present? YES Wetland hydrology present? YES	bed? (circle) atic? (circle) ap showing sampli	Are 'normal circumstal Explain answers in Rer ng point locations, tr	ransects, important features etc.
THE DOMESTIC AND THE STATE OF T	SECTION B	– VEGETATIC	DN .
Use scientific names of plants. Absolutive Tree Stratum (Plot size:	Species?	Indicator parture of Status Specific? ,,	Dominance Test:  No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  (B) 2  % OBL/FACW/FAC  (A/B) / OO  Prevalence Index:  Total % cover of:  Multiply by:  OBL
12. 207.= 27.17 Total cover = 135. Remarks:  * Kheed lox tuil - Lobels  Other species = JMV ant  - Small area an regetated	a perpunill		UNCERTAIN .



	SECT	TION A – S	SITE INFOR	MATION
Site: Homes Head Boy Owner: RCL groups Landform: Undulating land Is the land drained (circle) YES NO GPS (NZTM): E126 53 02  Are climatic/hydrologic conditions on Are vegetation, soil or hydrology significant in the soil of the soil or hydrology significant in the soil	Reg Dat Loc Inv N 49977	gion: Otago  e: 30/01/20  al relief: f(at estigator(s): Fni  +3 Altit  this time of year?	25 Gall ag hai ude m: 360 YES NO (cir	Sampling point: WA2 - plot 5  Land use: Cultivated & grazed  Land cover: excitic grassland  Soil °C: not measured slope°: n/a  Photo Nos: sare in report  Lolder R7483  rcle appropriate; if NO explain in Remarks)  umstances' present? (circle) YES NO
Are vegetation, soil or hydrology natu  SUMMARY OF FINDINGS—Att  Hydrophytic vegetation preser  Hydric soils present?  Wetland hydrology present?	ach site map s nt? YES YES YES	howing sampl	ing point locatio	in Remarks if needed  ns, transects, important features etc. a within a wetland? YES NO
Use scientific names of plants.  Tree Stratum (Plot size:	Absolute % cover — — — — — — — — — — — — — — — — — — —	Dominant Species?	FACW FACU FACU FACU FACU FACU	Dominance Test:  No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  % OBL/FACW/FAC  Prevalence Index:  Total % cover of:  OBL  FACW  FAC  FAC  UPL  Total  Morphological adaptations¹ (supporting data in Remarks)  Prevalence Index solisturbed or problematic  Hydrophytic vegetation present?  YES  NO  UNCERTAIN



		SECTIO	)N C -	SOIL AI	ND HYDROL	LOGY	
Profile descri	ption: (Describe to t	he depth needed to con	firm indicate	or presence/ab	sence, 30 cm default)		
Depth (cm)	Matrix colour (moist)	Mottles colour (moist) '	Mottles % <sup>1</sup>	Mottles Size <sup>2</sup>	Mottle location <sup>3</sup>	Material⁴	Remarks
0-17	6/3	5/8	3%	Medium	Matrix	mineral	
17-25	3/1	6/2	7%	Melium	Matrix	mineral	
25-32	6/2	5/2	77.	Medium	Mahix	mineral	
25-32	6/2	5/8	1%-	Fine	Matrix	mineral	
<sup>1</sup> Use % area c <b>Hydric soi</b> l Organic laye	indicators:	Ses; <sup>3</sup> Ped face, pore, wit	w mw 1(	P) VP	Cau	ty), humic, mineral soi use of wetness (circle action: Depression (Flat	appropriate):
	soil material	Iron concretions Manganese concre Nodular Consistence: Plastic Sticky Fluid	tions	Gley OR Mottled Horizon: Reductimor Redox mott Redox segre Perch-gley f	Wa Hig Par phic Par Lay egations Slov	ter table: Depth (cm) h GW (Perched) Seep is: Depth (cm)	
Hydric soils	present?	YES N	0	UNCE	RTAIN	NZSC subgroup	
Soil satu Water n Sedimer Drift dep  Gecondary Water-st Drainage Dry-seaso	water <30 cm (1B) rration <30 cm (1C) narks (2A) nt deposits (2B) nosits (2C) hydrology indica ained leaves (2K) patterns (2L) on water table (3E) n in aerial imagery (	Inundatio Sparsely v Salt crust  ators: minimum of Gec Sha	oil cracks (2F n on aerial in regetated co (2T)  2 require  morphic po flow aquitar  c-neutral tes	magery (2G) incave surface ( ed; check all isition (4B) ed (4C)	Oxidised r Reduced i Reduced i High wate  boxes that apply  FAC-neutral test ( 1. No. OBL & FAC	ron in tilled soil (3D) or table stunted/stress (4D); refer to Section E CW dominant species L dominant species	red plants (4A)
etland hy	drology present?	YES		NO			
100	soil:					Lean	v Ort
etch of site/s	The state of the s		<b>X</b> /	War.	worker	The way the plants of the plan	
marks:		Juntsd, Co	X/_	(()	uner	THE WAY THE PLOT	the state of the s



	SECT	TION A - S	SITE INFOR	MATION
Site: Homes Had Bay Owner: RCL Croup Landform: undulating land Is the land drained (circle) (YES) NO GPS (NZTM): £/265284 N499		cal relief: gradu	1025	gulles Landcover exotic availland
Are climatic/hydrologic conditions on th Are vegetation, soil or hydrology signific Are vegetation, soil or hydrology natural	antly disturbed	? (circle)	Are 'normal circ	rcle appropriate; if NO explain in Remarks) umstances' present? (circle) in Remarks if needed
Hydrophytic vegetation present?	YES YES YES	NO Is to	the sampled are	ens, transects, important features etc.  a within a wetland? YES NO
Use scientific names of plants.	Absolute	Dominant Dominant	– VEGETA	TION
Tree Stratum (Plot size:)  1. 2. 3. 4. Total cover = Sapling/Shrub Stratum (Plot size:	% cover	Species?	FAC W	Dominance Test:  No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  (B) 2  (B) 2  (B) 2  (B) 2  (B) 3  Prevalence Index:  Total % cover of:  Multiply by:  OBL 0
9				Hydrophytic vegetation present?  YES  NO  UNCERTAIN
Remarks: *Drain dug mithin co -No further tests			wired.	



## NEW ZEALAND WETLAND DELINEATION DATA FORM SECTION A – SITE INFORMATION Sampling point: WA2 - plot 7 Region: Otago Site: Homestead Bay Owner: RCL Group Date: 3/2/2025 Landform: Bently rolling hill country Local relief: flat Date: 3/2/2025 Land use: alltirated & grazed Local relief: flat Land cover: exotic grassland Soil °C: not recorded Slope : n/a Is the land drained (circle) YES NO Investigator(s): Erin Gallagher Altitude m: 363 GPS (NZTM): E126 5.288 N4997752 Photo Nos: saved in report folder Are climatic/hydrologic conditions on the site typical for this time of year? YES NO (circle appropriate; if NO explain in Remarks) Are vegetation, soil or hydrology significantly disturbed? (circle) Are 'normal circumstances' present? (circle) YES Are vegetation, soil or hydrology naturally problematic? (circle) Explain answers in Remarks if needed SUMMARY OF FINDINGS—Attach site map showing sampling point locations, transects, important features etc. Hydrophytic vegetation present? YES NO Is the sampled area within a wetland? YES No uncertain YES Hydric soils present? NO Wetland hydrology present? YES L SECTION B - VEGETATION Absolute Indicator Use scientific names of plants. Dominant Dominance Test: Tree Stratum (Plot size:\_\_\_\_) % cover Species? Status No. Dominant Spp. OBL/FACW/FAC (B) 3 Tot. Dominant Spp. across strata (A/B) <u>3</u>3 % OBL/FACW/FAC Prevalence Index: Total % cover of: Multiply by: Total cover = x1= 0 Sapling/Shrub Stratum (Plot size: FACU 6/ UPL Prevalence Index (B/A) = \_ Herb Stratum (Plot size: 2m<sup>2</sup>) Hydrophytic vegetation indicators: FACW 1. Juneus australis . Dominance Test is >50% FACU 2. Trifolium repens Prevalence Index is ≤3.0<sup>1</sup> 3. Agrostis capillaris FACU Morphological adaptations<sup>1</sup> (supporting data in 4. Leontodon saxatilis 1 Problematic hydrophytic vegetation<sup>1</sup> 5. Cynosurus instatus 6. Junais teniis <sup>1</sup>Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic 7. Lolium perenne Hydrophytic vegetation present? YES 11. UNCERTAIN V 20% = 23.8 Total cover = 119 50% = 59.5



Trome descript	ion: (Describe to th	e depth needed to cor	nfirm indicate	or presence/ab	sence, 30 cm default)		
Depth (cm)	Matrix colour (moist)	Mottles colour (moist)	Mottles %1	Mottles Size <sup>2</sup>	Mottle location <sup>3</sup>	Material <sup>4</sup>	Remarks
0-12	5/2	5/8	1	Median	Matrix	mineral	
12-18	4/2	6/3	15	Fine	Matrix	mineral	
18-28	6/3	5/2	15	Fine	Matrix	mineral	
28 - 32	4/2		-	1	-		Litter proots
¹Use % area ch	arts; <sup>2</sup> Use size class	es; <sup>3</sup> Ped face, pore, wit	thin ped alon	ng roots, within	matrix; <sup>4</sup> Organic (pea	ty), humic, minera	al soil
Hydric soil i	ndicators:	Soil drainage (circle)					rcle appropriate):
Organic layers		Concretions:		Colours: profile	form either.		Flat Valley Gully Slope
Organic so	oil material	Iron concretions  Manganese concre	etions [	Gley OR Mottled		ter table: Depth ( h GW (Perched):	cm) Seepage Tidal Lithic
Fibric		Nodular	L	Horizon:		s: Depth (cm ) _	
Mesic Humic	C	Consistence:		Reductimo	i pine		Densi- Duri- Fragi Ortstein
Peaty top:	soil	Plastic Sticky		Redox mot		ers: Depth (cm) _ w perm argillic	
Peaty sub	soil [	Fluid		Perch-gley		Pugged	
Hydric soils	present?	YES N	10	UNC	ERTAIN 📈	NZSC subgro	oup
Water ma Sediment Drift depo	t deposits (2B) osits (2C)	Inundation Sparsely Salt crust	vegetated co	imagery (2G) oncave surface ed; check al	(2H) Reduced i Reduced i High wate	ron in tilled soil (3 er table stunted/s	
	patterns (2L)		allow aquita			CW dominant spe	
	n water table (3E) n in aerial imagery (3		.C-neutral tes ost-heave hu	st (4D) immocks (4E)	2. No. FACU & UF 3. Total	L dominant speci	
					4. FAC-neutral (>	50%)	<u>33.3 (</u> A/A+B)*100
1	Irology present?	YES _		NO			
Wetland hyd	oil:						
Wetland hyd Sketch of site/so							
Sketch of site/so							
Sketch of site/so	11111111111111111111111111111111111111	1 Walst day	1 m 1/10	1 Wines	1sta wall than		
Sketch of site/so	William I want	1 1/11/1/1/1/1/1/1/1/1/1/1/1/1/1/1/1/1/	W 30, WW	1 Wester	1111 AND THE STREET		
Sketch of site/so	W. Herri	1 1/11/1/1/1/1/1/1/1/1/1/1/1/1/1/1/1/1/	W on WW.	1 Wyore	1010 May (1011) 18		



. INEVV ZEALAINE	VVLILAN	DELINEA	TION DATA FORM
SECT	ION A – S	ITE INFORM	MATION
Date and form: RCL Crowp Date country Local Steps (NZTM): E126 5398 N499784	Photo Nos: <u>Saved in report Folder</u> R 7483a		
Are climatic/hydrologic conditions on the site typical for Are vegetation, soil or hydrology significantly disturbed: Are vegetation, soil or hydrology naturally problematic?	? (circle)	Are 'normal circum	appropriate; if NO explain in Remarks) stances' present? (circle)  Remarks if needed
SUMMARY OF FINDINGS—Attach site map s Hydrophytic vegetation present? YES Hydric soils present? Wetland hydrology present?  YES  Wetland hydrology present?			i, transects, important features etc. within a wetland? YES NO
S	SECTION B	B – VEGETAT	ION
Use scientific names of plants. Absolute Tree Stratum (Plot size:) % cover 1 2	Dominant Species?	Indicator Status	Dominance Test:  No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  (B) /  % OBL/FACW/FAC  (A/B)
4			Total % cover of: Multiply by:  OBL $0$ $\times 1 = 0$ FACW $0$ $\times 2 = 0$ FAC $2$ $\times 3 = 6$ FACU $73.5$ $\times 4 = 294$ UPL $1$ $\times 5 = 5$ Total $77.5$ (A) $309$ (B)  Prevalence Index (B/A) = $3.99$
Herb Stratum (Plot size: 2m²)  1. Lolium perenne 70  2. Trifolium repens 1  3. Plantago australis 1  4. Leontodon saxatilis 1  5. Anthoanxthum odoratum 1  6. Verbascum virgatum 1  7. Cirsium valgane 2  8. Sagina porepens 0.5	<u>y</u> 	FACU FAC FAC FACU None→un FACU FACU	Dominance Test is >50%  Prevalence Index is ≤3.0¹  Morphological adaptations¹ (supporting data i Remarks)  Problematic hydrophytic vegetation¹  ¹Indicators of hydric soil and wetland hydrology mube present, unless disturbed or problematic
9			Hydrophytic vegetation present?  YES  NO  UNCERTAIN
Remarks:  No further tests requ	ured Juna	lertaken.	



S	ECTION A – S	SITE INFORMA	ATION
Site: Homestead Bay  Dwner: RCL Groups  Landform: gentle rolling hill country  s the land drained (circle) YES (NO  SPS (NZTM): F1265401 N4997	Date: 3/2/202 Local relief: con Con Investigator(s): Environmental Altitudes	me Gallagher ude m: 370	Sampling point: WA3-plot 9  Land use: cultivated & grazed  Land cover: exotic grass land.  Soil °C: not recorrectslope°: concare  Photo Nos: sared in report fold  R7483.  propriate; if NO explain in Remarks)
Are vegetation, soil or hydrology significantly dist	urbed? (circle)	Are 'normal circumsta	ances' present? (circle) YES NO
Are vegetation, soil or hydrology naturally probled SUMMARY OF FINDINGS—Attach site r		Explain answers in Re	
Hydrophytic vegetation present? YES Hydric soils present? YES Wetland hydrology present? YES	NO NO	ncortain - VEGETATIO	thin a wetland? YES  NO  NO
Use scientific names of plants.   Absolution	er Species?	Indicator Pasture \$ Status Species?   FAC FACU 9 0.1  None (UPL)  Total pasture = 0.1	Dominance Test:  No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  (B) 2  (A/B) 50%  Prevalence Index:  Total % cover of:  Multiply by:  OBL  FACW  FACW  FACW  O  FACW  O  FACU  O  I  FACU  O  I  FACU  O  I  FACU  O  FACU  O  FACU  O  FAC  FACU  O  FAC  I  FACU  O  FAC  I  FACU  O  FAC  I  FACU  O  I  Morphological adaptations¹ (supporting data in Remarks)  Problematic hydrophytic vegetation¹  Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic  Hydrophytic vegetation present?  YES  NO  UNCERTAIN
Remarks:  * likely FACU.  (P/TVL) × 100 = 3% pastine	species		



		SECTION	ON C -	SOIL A	ND HYDROL	.OGY	
Profile descri	ption: (Describe to	the depth needed to cor	firm indicate	or presence/ab	sence, 30 cm default)		
Depth (cm)	Matrix colour (moist)	Mottles colour (moist)	Mottles %1	Mottles Size <sup>2</sup>	Mottle location <sup>3</sup>	Material <sup>4</sup>	Remarks
0-21	6/1	6/8	5%	Medium	Matrix	mineral	
21-33	3/2	4/2	2%	Fine	Mamix	mireral	
	indicators:	Sses; <sup>3</sup> Ped face, pore, with Soil drainage (circle)	W MW I	P (VP)	Cau	use of wetness (cir	
Organic Litter Fibric	soil material	Iron concretions  Manganese concre  Nodular	etions [	Gley OR  Mottled  Horizon:	Wa Hig Par	ter table: Depth (c h GW Perched S ns: Depth (cm )	m) eepage Tidal Lithic 
Mesic Humic Peaty to Peaty su		Consistence: Plastic Sticky Fluid	.[	Reductimo Redox mot Redox segr Perch-gley	tled Lay regations Slo	Humus Fe-pan [ ers: Depth (cm) w perm argillic Pugged	Densi- Duri- Fragi Ortstein
Hydric soils	present?	YES N	10	UNC	ERTAIN	NZSC subgrou	מנ
Soil satu Water n	water <30 cm (1B) pration <30 cm (1C) parks (2A) pat deposits (2B) posits (2C)	Inundation	oil cracks (21 on on aerial i vegetated co	F) imagery (2G) oncave surface	Oxidised Reduced (2H) Reduced	sulphide odour (3 rhizosphere on roo iron (3C) iron in tilled soil (3 er table stunted/st	ots (3B)
econdary	hydrology indic	cators: minimum o	f <u>2</u> require	ed; check al	boxes that apply		
Drainage Dry-seas	ained leaves (2K) patterns (2L) on water table (3E) on in aerial imagery	Sh FA	omorphic po allow aquita C-neutral tes ost-heave hu	rd (4C)	1. No. OBL & FA	(4D); refer to Secti CW dominant specie PL dominant specie 50%)	ies(A)
Vetland hv	drology present	t? YES		NO	A STATE OF THE STA		
ketch of site/	NAME OF TAXABLE PARTY.		4 July 1	6 Ne 8 1935			
		25					



Are climatic/hydrologic conditions on the site typical for this time of year?  Are vegetation, soil or hydrology significantly disturbed? (circle)  Are vegetation, soil or hydrology naturally problematic? (circle)  Explain answers in Remarks if needed  SUMMARY OF FINDINGS—Attach site map showing sampling point locations, transects, important features etc.  Hydrophytic vegetation present? YES  NO  Is the sampled area within a wetland? YES  Hydric soils present?  YES  NO  SECTION B—VEGETATION  Use scientific names of plants.  Tree Stratum (Plot size:  No  Species?  Status  Species?  Total Cover =  Sapling/Shrub Stratum (Plot size:  1.  Total cover =  Sapling/Shrub Stratum (Plot size:  1.  2.  3.  4.  Total cover =  Sapling/Shrub Stratum (Plot size:  1.  2.  3.  4.  Total cover =  Sapling/Shrub Stratum (Plot size:  1.  2.  3.  4.  Total cover =  Sapling/Shrub Stratum (Plot size:  1.  2.  3.  4.  Total cover =  Sapling/Shrub Stratum (Plot size:  1.  2.  3.  4.  Total cover =  Sapling/Shrub Stratum (Plot size:  1.  2.  3.  4.  Total cover =  Sapling/Shrub Stratum (Plot size:  1.  2.  3.  4.  Total cover =  Sapling/Shrub Stratum (Plot size:  1.  2.  3.  4.  Total cover =  Sapling/Shrub Stratum (Plot size:  1.  2.  3.  4.  5.  Total cover of:  FACU  FAC	nzed on care t bolde R 7483
Are vegetation, soil or hydrology significantly disturbed? (circle)  Are vegetation, soil or hydrology naturally problematic? (circle)  SUMMARY OF FINDINGS—Attach site map showing sampling point locations, transects, important features etc.  Hydrophytic vegetation present? YES NO Is the sampled area within a wetland? YES NO  Wetland hydrology present? YES NO NO  SECTION B — VEGETATION  Use scientific names of plants. Absolute Dominant Tree Stratum (Plot size: % cover Species? Status Species? Status Species? Status Species? No Dominant Spp. OBL/FACW/FAC Tot. Dominant Spp. across strata % OBL/FACW/FAC (A)  Total cover = Sapling/Shrub Stratum (Plot size: 10 Total cover = 10 Total cover of: Multiply by: 10 Total cover of: Multiply by: 10 Total cover of:	
SUMMARY OF FINDINGS—Attach site map showing sampling point locations, transects, important features etc.  Hydrophytic vegetation present? YES NO Is the sampled area within a wetland? YES Hydric soils present?  Wetland hydrology present? YES NO NO  SECTION B — VEGETATION  Use scientific names of plants.  Tree Stratum (Plot size:	(A) /
Use scientific names of plants. Absolute Dominant Indicator fashers \$\frac{1}{2}\$ Dominance Test:  Tree Stratum (Plot size:) % cover Species? Status species? Status species? No. Dominant Spp. OBL/FACW/FAC Tot. Dominant Spp. across strata % OBL/FACW/FAC (A/2)  Total cover = Sapling/Shrub Stratum (Plot size:)  Total cover = Sapling/Shrub Stratum (Plot size:)  Sapling/Shrub Stratum (Plot size:)  FACU	(A) /
Tree Stratum (Plot size:) % cover	(A) /
Total cover =  Herb Stratum (Plot size: $2m^2$ )  1. Tirbolium regens 2. Glycnia Pluitans 3. Solanum nigram 4. Agyortis capillaris 5. Duphania punilio 6. 7. 8. 9. 10. 11. 12. 20% = 0.68 Total cover = 3.4 Total cover = 3.4 Total parture FOUR - 1.7 Total 3.4 (A) 7.7 (E  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index (B/A) = 2.26  Nore (UR)  UNCERTAIN UNCERTAIN UNCERTAIN UNCERTAIN UNCERTAIN UNCERTAIN UNCERTAIN UNCERTAIN UNCERTAIN UNCERTAI	(B) Z



	iption: (Describe to th	ne depth needed to cor	firm indicate	or presence/al	sence 30 cm	default)	
Depth (cm)	Matrix colour (moist)	Mottles colour (moist)	Mottles %1	Mottles Size <sup>2</sup>	Mottle loc		Remarks
0-17	6/1	5/8	5	Fine	Matrix	mireral	
17-30	3/2	6/2	1	Fine	Matri	x miheral	
¹Use % area	charts; <sup>2</sup> Use size class	es: <sup>3</sup> Ped face, pore, wit	hin ned alon	og roots within	matrix: <sup>4</sup> Orga	nic (peaty), humic, mine	ural coil
Hydric so	il indicators:	Soil drainage (circle)		-	illiatrix, Orga		circle appropriate):
Organic lay	ers:	Concretions:		Colours: profile	form either:	Location: Depression	on Flat Valley Gully Slope
=	soil material	Iron concretions		Gley OR		Water table: Depth	
Litter		Manganese concre		Mottled		High GW Perched Pans: Depth (cm ) _	Seepage Tidal Lithic
Mesic		Consistence:	-	Horizon:	orphic		Densi- Duri- Fragi Ortste
Humic		Plastic		Redox mo		Layers: Depth (cm)	
Peaty t		Sticky		Redox seg		Slow perm argillic	
		Fluid		Perch-gley	features	Pugged	
Hydric soil	s present?	YES /	10	UNC	ERTAIN	NZSC subgr	oup
Primary	hydrology indic	ators: minimum o	f <u>1</u> require	ed; check all	boxes that	apply	
Surface	e water (1A)	Algal ma	t/crust (2D)		□ A	quatic invertebrates (2J)	
	dwater <30 cm (1B)	Iron depo			□ <sup>+</sup>	lydrogen sulphide odour	(3A)
	curation <30 cm (1C) marks (2A)		oil cracks (2f	magery (2G)		oxidised rhizosphere on r	oots (3B)
	ent deposits (2B)			ncave surface	/	educed iron (3C) educed iron in tilled soil	(3D)
Drift de	eposits (2C)	Salt crust	(21)			ligh water table stunted/	
Secondary	hydrology indic	ators: minimum o	f <u>2</u> require	ed; check al	l boxes that	apply	
Water-	stained leaves (2K)	Ge	eomorphic po	osition (4B)	FAC-neut	tral test (4D); refer to Se	ction B: Vegetation
	ge patterns (2L)	THE RESERVE TO SERVE THE PARTY OF THE PARTY	allow aquita			BL & FACW dominant sp	
=	son water table (3E)		C-neutral tes			CU & UPL dominant spe	cies / (B) _ Z (A+B)
Saturat	ion in aerial imagery	(3F)	ost-neave nu	mmocks (4E)	3. Total 4. FAC-ne	eutral (>50%)	50 (A/A+B)*:
Motland b	ydrology present	? YES 🗸		NC			
Sketch of site		: 153 [V]		NC			
	mindle adding					nd.	,
	A	were f	PIO			west	
		The.		L. Y	Walter	Willes and the second	
		4	1	110			
Remarks:						storksbill, A	



NEW ZI				TION DATA FORM
	SECTION	S – A NC	ITE INFORM	
Owner: RCL Groups  Landform: Partle rolling hill course  Is the land drained (circle) YES (NO  GPS (NZTM): F126 5634 N4	Soil C: Not recorded Slope			
Are climatic/hydrologic conditions on the Are vegetation, soil or hydrology significa Are vegetation, soil or hydrology naturall	ntly disturbed? (o	circle) ircle)	Are 'normal circun  Explain answers in	e appropriate; if NO explain in Remarks) estances' present? (circle)  Remarks if needed
SUMMARY OF FINDINGS—Attac Hydrophytic vegetation present? Hydric soils present? Wetland hydrology present?	YES TO YES	NO Is t	ng point location: the sampled area	s, transects, important features etc. within a wetland? YES NO NO
Use scientific names of plants.  Free Stratum (Plot size:)  12.	Absolute % cover	Dominant Species?	Indicator Status	Dominance Test:  No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  % OBL/FACW/FAC  (A/B)
3				Prevalence Index:  Total % cover of:  OBL  OBL  FACW  Total  FACU  FACU  Multiply by: $0$ $0$ $0$ $0$ $0$ $0$ $0$ $0$
Herb Stratum (Plot size: 2m²)  1. Cirsium rulgare  2. Cirsium arrense  3. Chens podium album  4. Lolium percune  5. Trifolium repens  6. Dysphania pumilio  7. Solanum nigram  8. Urtica urens	10	<u>y</u> <u>y</u>	FACU None FACU FACU None FACU None	Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index is ≤3.0¹  Morphological adaptations¹ (supporting data in Remarks)  Problematic hydrophytic vegetation¹  ¹Indicators of hydric soil and wetland hydrology musbe present, unless disturbed or problematic
9. Verbrocum Mapsus 10. 10a annua 11. Plantago australia 12. 201-= 4.1 Total cover= 501-= 10.25	0.5	<u>y</u>	Nore FACU FAU	Hydrophytic vegetation present? YES
Remarks:  No further assess,	nents r	equired /	'under tak	ln.



	SECT	ION A - S	ITE INFOR	MATION
Is the land during of (single) VES (NO)		e: 3/2/202 al relief: <u>con cu</u> estigator(s): <u>E min</u> Altitu	ude m: 362	Land use: grazed pasture
Are vegetation, soil or hydrology significant Are vegetation, soil or hydrology naturally				mstances' present? (circle) YES NO n Remarks if needed
Hydrophytic vegetation present? \ Hydric soils present?	YES	NO Is		ns, transects, important features etc. a within a wetland? YES NO
Use scientific names of plants.	Absolute	Dominant	Indicator	Dominance Test:
Tree Stratum (Plot size:)  1	1 	Species?	FACU FACU FACU FACU FACU Vone None	No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  (B) /  W OBL/FACW/FAC  (A/B) \( \text{N} \)  Prevalence Index:  Total % cover of: Multiply by:  OBL \( \frac{0}{2} \) \( \text{x1} = \frac{0}{2} \)  FACW \( \frac{0}{2} \) \( \text{x3} = \frac{0}{2} \)  FACU \( \frac{6}{2} \) \( \text{x4} = \frac{24}{24} \)  UPL \( \frac{0}{2} \) \( \text{x5} = \frac{0}{2} \)  Total \( \frac{6}{2} \) \( \text{(A)} \) \( \text{24} \) \( \text{(B)} \)  Prevalence Index (B/A) = \( \frac{4}{2} \) \( \text{0} \)  Hydrophytic vegetation indicators:  \[ \text{Dominance Test is >50%} \]  Prevalence Index is \( \frac{3}{2} \).  Morphological adaptations\( ^1 \) (supporting data in Remarks)  \[ \text{Problematic hydrophytic vegetation} ^1  \]  \[ \text{1-indicators of hydric soil and wetland hydrology mus be present, unless disturbed or problematic}
8.				Hydrophytic vegetation present? YES
50°10: 3.35 Remarks: - No Curther RSS E	isments	require	d/underta	ken.



	SECTION	A - SITE IN	FORMATION	
Site: Homestead Bay Owner: RCL Group Landform: Jentu rolling will war Is the land drained (circle) YES (NO) GPS (NZTM): £126,5442 N	Date: 4/.  Local relief: Investigator	2/2025 slightly con c (s): Erin Gallagh	Land  Land  Land  Soil of	ling point: WAS- plot /3 use: authinated 3 grazed. cover: exotic grass land c: not recorded slope: flat a Nos: saved in report fol. R78
Are climatic/hydrologic conditions on the Are vegetation, soil or hydrology significa Are vegetation, soil or hydrology naturall	ntly disturbed? (circle)	Are 'nor	NO (circle appropriate mal circumstances' presonances in Remarks if no	
SUMMARY OF FINDINGS—Attac Hydrophytic vegetation present? Hydric soils present? Wetland hydrology present?	YES N		ed area within a w	
Use scientific names of plants. Tree Stratum (Plot size:) 1 2 3	Absolute Do		Pasture S Domina Species? : No. Do Tot. Do	minant Spp. OBL/FACW/FAC (A)
			Total % OBL FACW FAC FACU UPL Total	2 x4= 8
3. Johin perence 4. Malra reglecta 5. Cirsiam vulgare 6. Tribohin repens 7. Conium manulatum	2 / 0.1 0.5 0.5 0.5	FAC	£)	ohytic vegetation indicators:  ominance Test is >50%  evalence Index is ≤3.0¹  orphological adaptations¹ (supporting darmarks)  oblematic hydrophytic vegetation¹  tors of hydric soil and wetland hydrology sent, unless disturbed or problematic
8. Rumex obt 9. 10. 11. 12. 20% - 6.94 Total cover = 50% > 17.35	34.7	Total pasture species cover	Hydro YES NO UNCE	pphytic vegetation present?
Remarks:  Other species Learby  No further asses	1 = MElalp, R	Osab, VERVIN	Hordeum g	



are vegetation, soil or hydrology significantly disturbed? (circle)  Are 'normal circums	Land cover: exotic grasslard  Soil °C: not recorded Slope : concare, to
SUMMARY OF FINDINGS—Attach site map showing sampling point locations,	nstances' present? (circle)  Remarks if needed  The transects, important features etc.
Hydric soils present?  YES NO is the sampled area with the sampled	NO NO NO
Se scientific names of plants.  Absolute  Dominant  Indicator  Free Stratum (Plot size:)  Total cover =	Dominance Test:  No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  % OBL/FACW/FAC  Prevalence Index:  Total % cover of:  Multiply by:  OBL  FACW  O  FACW  O  FAC  UPL  O  Total  O  (A)  Prevalence Index  Total  FACW  O  FAC  O  O  FAC  O



Profile descrip	otion: (Describe to th	he depth needed to cor	nfirm indicate	or presence/ab	sence, 30 cm default)		
Depth (cm)	Matrix colour (moist)	Mottles colour (moist)	Mottles %1	Mottles Size <sup>2</sup>	Mottle location <sup>3</sup>	Material <sup>4</sup>	Remarks
0-12	4/3	4/6	1	Fine	Matrix	mineral	stong layer, hard dig through.
12-30	3/2	2/2	5%.	Medium	Mahix	mineral	
	harts; <sup>2</sup> Use size class I indicators:	ses; <sup>3</sup> Ped face, pore, wi	thin ped alo	ng roots, withir			
Organic laye		Soil drainage (circle)  Concretions:  Iron concretions  Manganese concr  Nodular	etions	Colours: profile Gley OR Mottled Horizon:	e form either: Wi	cation: Depressionater table: Depth	ircle appropriate):  The Flat Valley Gully Slope  (cm)  Seepage Tidal Lithic
Mesic Humic Peaty to		Consistence:  Plastic Mergin Sticky Fluid	nd	Reductimo Redox mo Redox seg Perch-gley	ttled Lay regations Slo	vers: Depth (cm) _	Densi- Duri- Fragi Ortstein  Ly at lowest point
		VEC .	NO	l unc	CERTAIN T	N7CC subare	
Primary  Surface  Ground	s present?  hydrology indice  water (1A)  dwater <30 cm (1B)  turation <30 cm (1C)	cators: minimum o		ed; check al	I boxes that apply Aquatic i Hydroger	NZSC subgro	(3A)
Primary Surface Ground Soil sat Water Sedime	hydrology indice water (1A) dwater <30 cm (1B)	Algal man liron dep	of 1 requir at/crust (2D) posits (2E) soil cracks (2 tion on aerial y vegetated of	ed; check al	Aquatic i Hydrogei Oxidised Reduced Reduced	nvertebrates (2J) n sulphide odour rhizosphere on ro iron (3C) iron in tilled soil (	(3A) pots (3B)
Primary Surface Ground Soil sai Water Sedim Drift d	r hydrology indice water (1A) dwater <30 cm (1B) turation <30 cm (1C) marks (2A) ent deposits (2B) eposits (2C)	Algal man department of the surface	of 1 requires (2D) posits (2E) soil cracks (2 cition on aerial by vegetated of st (2I)	red; check all  2F) I imagery (2G) concave surface	Aquatic i Hydroger Oxidised Reduced Reduced High wat	nvertebrates (2J) n sulphide odour i rhizosphere on ro iron (3C) iron in tilled soil ( er table stunted/s	(3A) pots (3B)
Primary Surface Ground Soil sai Water Sedim Drift d  Secondare Draina Dry-see Satura	whydrology indice water (1A) dwater <30 cm (1B) turation <30 cm (1C) marks (2A) ent deposits (2B) eposits (2C)  y hydrology indices stained leaves (2K) ge patterns (2L) ason water table (3E tion in aerial imagen	Algal ma Iron dep Iro	of 1 requires at/crust (2D) posits (2E) soil cracks (2 ion on aerial y vegetated cost (2T)  of 2 requires accomplic phallow aquit AC-neutral to	ed; check all  2F)  I imagery (2G)  concave surface  red; check all  possition (4B)  ard (4C)	Aquatic i Hydroger Oxidised Reduced High wat  II boxes that apply  FAC-neutral test 1. No. OBL & FA 2. No. FACU & U 3. Total	nvertebrates (2J) n sulphide odour of thizosphere on rolling iron (3C) iron in tilled soil (er table stunted/sulphi); refer to Sec. CW dominant spec	(3A)  bots (3B)  (3D)  stressed plants (4A)  tion B: Vegetation exies(A) exies(B)(A+B)
Primary Surface Ground Soil sai Water Sedim Drift d  Secondary  Water- Draina Dry-see Satura	whydrology indice water (1A) dwater <30 cm (1B) turation <30 cm (1C) marks (2A) ent deposits (2B) eposits (2C)  y hydrology indi estained leaves (2K) ge patterns (2L) ason water table (3E)	Algal ma Surface Inundat Sparsely Salt crusticators: minimum of Surface Inundat Sparsely Salt crusticators: minimum of Sparsely Salt cr	of 1 requires at/crust (2D) posits (2E) soil cracks (2 ion on aerial y vegetated cost (2T)  of 2 requires accomplic phallow aquit AC-neutral to	red; check all limagery (2G) concave surface red; check all position (4B) ard (4C) est (4D) nummocks (4E)	Aquatic i Hydroger Oxidised Reduced High wat Il boxes that apply FAC-neutral test 1. No. OBL & FA 2. No. FACU & U 3. Total 4. FAC-neutral (s	nvertebrates (2J) n sulphide odour of thizosphere on rolling iron (3C) iron in tilled soil (er table stunted/sulphi); refer to Sec. CW dominant spec	(3A)  pots (3B)  (3D)  stressed plants (4A)  tion B: Vegetation exies(A) elies(B)
Primary  Surface Ground Soil sar Water Sedim Drift d  Secondar  Water Draina Dry-se Satura Google tax	whydrology indice water (1A) dwater <30 cm (1B) turation <30 cm (1C) marks (2A) ent deposits (2B) eposits (2C) whydrology indicestained leaves (2K) ge patterns (2L) ason water table (3E tion in aerial imagent of the control of the	Algal ma Surface Inundate Sparsely Salt crusticators: minimum of Surface Inundate Sparsely Salt crusticators: minimum of Sparsely Sparsely Salt crusticators: minimum of Sparsely Spars	of 1 requires at/crust (2D) posits (2E) soil cracks (2 ion on aerial y vegetated cost (2T)  of 2 requires accomplic phallow aquit AC-neutral to	ed; check all limagery (2G) concave surface red; check al possition (4B) ard (4C) est (4D)	Aquatic i Hydroger Oxidised Reduced High wat Il boxes that apply FAC-neutral test 1. No. OBL & FA 2. No. FACU & U 3. Total 4. FAC-neutral (s	nvertebrates (2J) n sulphide odour of thizosphere on rolling iron (3C) iron in tilled soil (er table stunted/sulphi); refer to Sec. CW dominant spec	(3A)  bots (3B)  (3D)  stressed plants (4A)  tion B: Vegetation exies(A) exies(B)(A+B)
Primary  Surface Ground Soil sa Water Sedim Drift d  Secondar  Water Draina Dry-se Satura Google too	whydrology indice water (1A) dwater <30 cm (1B) turation <30 cm (1C) marks (2A) ent deposits (2B) eposits (2C)  y hydrology indices water table (3E) ason water table (3E) tion in aerial imagen  who 11/2018, 3/2 mydrology preser e/soil:	Algal ma Surface Inundat Sparsely Salt crusticators: minimum of Sparsely Salt crusticators: minimum of Sparsely Salt crusticators: minimum of Sparsely The Sparsely Salt crusticators: minimum of Sparsely The Sparsely Spa	of 1 requires at/crust (2D) posits (2E) soil cracks (2 ion on aerial y vegetated cost (2T)  of 2 requires accomplic phallow aquit AC-neutral to	red; check all limagery (2G) concave surface red; check all position (4B) ard (4C) est (4D) nummocks (4E)	Aquatic i Hydrogei Oxidised Reduced High wat  Il boxes that apply  FAC-neutral test 1. No. OBL & FA 2. No. FACU & U 3. Total 4. FAC-neutral (3	nvertebrates (2J) n sulphide odour rhizosphere on ro iron (3C) iron in tilled soil ( er table stunted/s  (4D); refer to Sec CW dominant spec PL dominant spec	(3A) pots (3B)  stressed plants (4A)  tion B: Vegetation ecies(A) cies(B)(A+B)(A/A+B)*100
Primary  Surface Ground Soil sa Water Sedim Drift d  Secondar  Water Draina Dry-se Satura Google too	whydrology indice water (1A) dwater <30 cm (1B) turation <30 cm (1C) marks (2A) ent deposits (2B) eposits (2C) whydrology indicestained leaves (2K) ge patterns (2L) ason water table (3E tion in aerial imagent of the control of the	Algal ma Surface Inundat Sparsely Salt crusticators: minimum of Sparsely Salt crusticators: minimum of Sparsely Salt crusticators: minimum of Sparsely The Sparsely Salt crusticators: minimum of Sparsely The Sparsely Spa	of 1 requires at/crust (2D) posits (2E) soil cracks (2 ion on aerial y vegetated cost (2T)  of 2 requires accomplic phallow aquit AC-neutral to	red; check all limagery (2G) concave surface red; check all position (4B) ard (4C) est (4D) nummocks (4E)	Aquatic i Hydrogei Oxidised Reduced High wat  Il boxes that apply  FAC-neutral test 1. No. OBL & FA 2. No. FACU & U 3. Total 4. FAC-neutral (3	nvertebrates (2J) n sulphide odour rhizosphere on ro iron (3C) iron in tilled soil ( er table stunted/s  (4D); refer to Sec CW dominant spec PL dominant spec	(3A) pots (3B)  stressed plants (4A)  tion B: Vegetation ecies(A) cies(B)(A+B)(A/A+B)*100
Primary  Surface Ground Soil sa Water Sedim Drift d  Secondar  Water Draina Dry-se Satura Google too	whydrology indice water (1A) dwater <30 cm (1B) turation <30 cm (1C) marks (2A) ent deposits (2B) eposits (2C)  y hydrology indices water table (3E) ason water table (3E) tion in aerial imagen  who 11/2018, 3/2 mydrology preser e/soil:	Algal ma Surface Inundat Sparsely Salt crusticators: minimum of Sparsely Salt crusticators: minimum of Sparsely Salt crusticators: minimum of Sparsely The Sparsely Salt crusticators: minimum of Sparsely The Sparsely Spa	of 1 requires at/crust (2D) posits (2E) soil cracks (2 ion on aerial y vegetated of st (2T) of 2 requires a requirement of 2 requires a restriction aquit AC-neutral to the rost-heave heave heave the statement of the rost-heave heave the restriction at the rost-heave heave the restriction at the restriction and the rost-heave heave the restriction at the	red; check all limagery (2G) concave surface red; check all position (4B) ard (4C) est (4D) nummocks (4E)	Aquatic i Hydroger Oxidised Reduced High wat Il boxes that apply FAC-neutral test 1. No. OBL & FA 2. No. FACU & U 3. Total 4. FAC-neutral (s	nvertebrates (2J) n sulphide odour rhizosphere on ro iron (3C) iron in tilled soil ( er table stunted/s  (4D); refer to Sec CW dominant spec PL dominant spec	(3A) pots (3B)  stressed plants (4A)  tion B: Vegetation ecies(A) cies(B)(A+B)(A/A+B)*100



NEW ZEAL	AND WETLAN	ID DELINEA	TION DATA FORM
	ECTION A – S		
Site: Homeskad Bay  Owner: RCL Group  Landform: Gentle relling win try  Is the land drained (circle) YES NO  GPS (NZTM): £1265267 N49981	Region: Otago  Date: 4/2/2025  Local relief: depute Investigator(s): Frir	5 spin r Gallagher ude m:_381	Sampling point: WAG- plot 15  Land use: grazed pasture  Land cover: arotic grassland  Soll °C: not regrated Slope°: flat francas  Photo Nos: saved wto report  folder R748.
Are climatic/hydrologic conditions on the site type Are vegetation, soil or hydrology significantly dis Are vegetation, soil or hydrology naturally proble SUMMARY OF FINDINGS—Attach site Hydrophytic vegetation present? YES Hydric soils present?	oical for this time of year? turbed? (circle) ematic? (circle) map showing sampli	YES NO (circle Are 'normal circum Explain answers in I	appropriate; if NO explain in Remarks) stances' present? (circle) YES NO Remarks if needed
Wetland hydrology present? YES	□ □ NO	VECETAT	
Use scientific names of plants. Abso		- VEGETAT	Dominance Test:
Tree Stratum (Plot size:)	over Species?	Status	No. Dominant Spp. OBL/FACW/FAC  Tot. Dominant Spp. across strata  % OBL/FACW/FAC  Prevalence Index:  Total % cover of: Multiply by:
Total cover =		=	OBL $0$ $x1 = 0$ FACW $0$ $x2 = 0$ FAC $0$ $x3 = 0$ FACU $3$ $x4 = 12$ UPL $3.5$ $x5 = 17.5$ Total $6.5$ (A) $29.5$ (B) Prevalence Index (B/A) = $4.54$
5. Solanum nigrum 0.		FACU FACU FACU FACU None (UPL)	Hydrophytic vegetation indicators:  Dominance Test is >50%  Prevalence Index is ≤3.0¹  Morphological adaptations¹ (supporting data in Remarks)  Problematic hydrophytic vegetation¹  ¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic
8			Hydrophytic vegetation present? YES
Remarks: No further assessme	nts undet	iakin /requ	ired.



## NEW ZEALAND WETLAND DELINEATION DATA FORM SECTION A - SITE INFORMATION Site: Homestead Bay Region: Otago Sampling point: Plot 16 Owner: RCL Homestead Bay Ltd Date: 3 September 2025 Land use: Farmland - cattle grazing Landform: Gentle undulating toeslope Local relief: Flat to mild slope Land cover: Exotic grassland Is the land drained (circle) YES NO Investigator(s): Erin Gallagher Soil °C: Not measured Slope°: Not measured GPS (NZTM): E1265457 N4998897 Altitude m: 363 m Photo Nos: Saved in Report folder 7522 Are climatic/hydrologic conditions on the site typical for this time of year? YES NO (circle appropriate; if NO explain in Remarks) Are vegetation, soil or hydrology significantly disturbed? (circle) Are 'normal circumstances' present? (circle) Are vegetation, soil or hydrology naturally problematic? (circle) Explain answers in Remarks if needed SUMMARY OF FINDINGS—Attach site map showing sampling point locations, transects, important features etc. Hydrophytic vegetation present? YES NO Is the sampled area within a wetland? YES V NO Hydric soils present? YES NO YES 🗸 Wetland hydrology present? SECTION B - VEGETATION Indicator Use scientific names of plants. Absolute Dominant Dominance Test: Tree Stratum (Plot size:\_\_\_\_ % cover Species? Status No. Dominant Spp. OBL/FACW/FAC (A) 1 Tot. Dominant Spp. across strata (B) 1 (A/B) 100% % OBL/FACW/FAC Prevalence Index: Total % cover of: Multiply by: Total cover = OBL 0 x1=0 Sapling/Shrub Stratum (Plot size:\_\_\_\_ 2 x2=4 FACW x3=294 FAC 98 x4=11.2 FACU 2.8 UPL 0 102.8 (A) Prevalence Index (B/A) = 3.1Total cover = Herb Stratum (Plot size: 2m x 2m) Hydrophytic vegetation indicators: 1. Holcus lanatus Dominance Test is >50% 2. Agrostis stolonifera FACW Prevalence Index is ≤3.01 3. Trifolium repens FACU Morphological adaptations1 (supporting data in 4. Lolium perenne Problematic hydrophytic vegetation<sup>1</sup> 5. Achillea millefolium 6. Agrostis capillaris <sup>1</sup>Indicators of hydric soil and wetland hydrology must 7. Galium aparine be present, unless disturbed or problematic Hydrophytic vegetation present? YES 10. NO UNCERTAIN V 102.8 Remarks:



## SECTION C - SOIL AND HYDROLOGY Profile description: (Describe to the depth needed to confirm indicator presence/absence, 30 cm default) Depth (cm) Matrix colour Mottles colour Mottles Mottles Mottle location<sup>3</sup> Material<sup>4</sup> Remarks 961 Size (moist) (moist) 0-40cm 4/1 op 10cm with grass roots/debri 5/1 2% Fine Matrix use % area charts; <sup>2</sup>Use size classes; <sup>3</sup>Ped face, pore, within ped along roots, within matrix; <sup>4</sup>Organic (peaty), humic, mineral soil Hydric soil indicators: Cause of wetness (circle appropriate): Soil drainage (circle) W MW I P VP Location: Depression Flat Valley Gully Slope Organic layers: Concretions Colours: profile form either: Water table: Depth (cm) Organic soil material Iron concretions Gley profile form OR High GW Perched Seepage Tidal Lithic Litter Manganese concretions Mottled profile form Pans: Depth (cm ) \_ Fibric Nodular Pan Humus Fe-pan Densi- Duri- Fragi Ortstein Mesic Consistence: Reductimorphic Restricting layer: Depth (cm) \_ Humic Plastic Redox mottled Slow perm argillic Peaty topsoil Sticky Redox segregations Peaty subsoil Fluid Pugged Perch-gley features Hydric soils present? YES UNCERTAIN | NZSC subgroup Primary hydrology indicators: minimum of 1 required; check all boxes that apply Algal mat/crust (2D) Surface water (1A) Aquatic invertebrates (2J) Groundwater <30 cm (1B) Iron deposits (2E) Hydrogen sulphide odour (3A) Surface soil cracks (2F) Soil saturation <30 cm (1C) Oxidised rhizosphere on roots (3B) Inundation on aerial imagery (2G) Water marks (2A) Reduced iron (3C) Sediment deposits (2B) Sparsely vegetated concave surface (2H) Reduced iron in tilled soil (3D) Drift deposits (2C) Salt crust (21) High water table stunted/stressed plants (4A) Secondary hydrology indicators: minimum of 2 required; check all boxes that apply FAC-neutral test (4D); refer to Section B: Vegetation Water-stained leaves (2K) Geomorphic position (4B) 1. No. OBL & FACW dominant species Drainage patterns (2L) Shallow aquitard (4C) Dry-season water table (3E) FAC-neutral test (4D) 2. No. FACU & UPL dominant species (B) Saturation in aerial imagery (3F) Frost-heave hummocks (4E) (A+B) 0 4. FAC-neutral (>50%) (A/A+B)\*100 YES V Wetland hydrology present? NO Sketch of site/soil: Remarks:

Call Free 0508 WILDNZ Ph +64 7 343 9017 Fax +64 7 349018 ecology@wildlands.co.nz

99 Sala Street PO Box 7137, Te Ngae Rotorua 3042, New Zealand

Regional Offices located in Auckland; Christchurch; Dunedin; Hamilton; Invercargill; Queenstown; Tauranga; Wānaka; Wellington; Whangārei.

